

**UNITED STATES PATENT APPLICATION**

**OF**

**URSULA SCHINDLER, KARL SCHÖNAFINGER, and HARTMUT STROBEL**

**FOR**


**SULFUR SUBSTITUTED SULFONYLAMINOCARBOXYLIC ACID N-ARYLAMIDES,  
THEIR PREPARATION, THEIR USE AND PHARMACEUTICAL PREPARATIONS  
COMPRISING THEM**

Amend.  
B<sup>1</sup>

The present application claims priority under 35 U.S.C. § 119 to German patent applications No. 19830430.7 filed July 8, 1998, and No. 19903126.6 filed January 27, 1999. Both priority applications are entirely incorporated herein by reference.

5

The present invention relates to compounds of the formula I



15 wherein A<sup>1</sup>, A<sup>2</sup>, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, X and n are as defined below, which are valuable  
pharmaceutically active compounds for the therapy and prophylaxis of diseases, for  
example for cardiovascular diseases such as hypertension, angina pectoris, cardiac  
insufficiency, thromboses or atherosclerosis. The compounds of the formula I are  
capable of modulating the body's production of cyclic guanosine monophosphate  
20 ("cGMP") and are generally suitable for the therapy and prophylaxis of diseases which  
are associated with a disturbed cGMP balance. The invention furthermore relates to

processes for preparing compounds of the formula I, to their use for the therapy and prophylaxis of the abovementioned diseases and for preparing pharmaceuticals for this purpose, and to pharmaceutical preparations which comprise compounds of the formula I.

5

cGMP is an important intracellular messenger which triggers a multitude of different effects via the modulation of cGMP-dependent protein kinases, phosphodiesterases and ion channels. Examples are the relaxation of smooth muscles, the inhibition of thrombocyte activation and the inhibition of the proliferation of smooth-muscle cells and of leukocyte adhesion. cGMP is produced by particulate and soluble guanylate cyclases as a response to a number of extracellular and intracellular stimuli. In the case of the particulate guanylate cyclases, stimulation is essentially effected by peptidic messengers, such as the atrial natriuretic peptide or the cerebral natriuretic peptide. The soluble guanylate cyclases ("sGC"), which are cytosolic heterodimeric heme proteins, in contrast, are essentially regulated by a family of low-molecular-weight factors which are formed enzymatically. The most important stimulant is nitrogen monoxide ("NO") or a closely related species. The function of other factors such as carbon monoxide or the hydroxyl radical is still largely unclear. The binding of NO to the heme with formation of a penta-coordinate heme-nitrosyl complex is being discussed as activation mechanism of the activation by NO. The associated release of the histidine which is bound in the basal state to the iron converts the enzyme into the active conformation.

Active soluble guanylate cyclases are composed of an  $\alpha$  and a  $\beta$  subunit each. Several subunit subtypes have been described which differ from one another with respect to sequence, tissue-specific distribution and expression in different development stages.

The subtypes  $\alpha_1$  and  $\beta_1$  are mainly expressed in brain and lung, while  $\beta_2$  is found in particular in liver and kidney. The subtype  $\alpha_2$  was shown to be present in human fetal brain. The subunits referred to as  $\alpha_3$  and  $\beta_3$  were isolated from human brain and are homologous to  $\alpha_1$  and  $\beta_1$ . More recent works indicate an  $\alpha_{2i}$  subunit which contains an insert in the catalytic domain. All subunits show great homologies in the region of the catalytic domain. The enzymes presumably contain one heme per heterodimer, which is bound via  $\beta_1$ -Cys-78 and/or  $\beta_1$ -His-105 and is part of the regulatory center.

Under pathologic conditions, the formation of guanylate-cyclase-activating factors can be reduced, or their degradation may be promoted owing to the increased occurrence of free radicals. The resulting reduced activation of the sGC leads, via a weakening of the respective cGMP-mediated cellular response, for example to an increase of the blood pressure, to platelet activation or to increased cell proliferation and cell adhesion. As a consequence, formation of endothelial dysfunction, atherosclerosis, hypertension, stable or unstable angina pectoris, thromboses, myocardial infarction, strokes or erectile dysfunction results. Pharmacological stimulation of sGC offers a possibility to normalize

cGMP production and therefore makes possible the treatment and/or prevention of such disorders.

For the pharmacological stimulation of the sGC, use has hitherto almost exclusively

5 been made of compounds whose activity is based on an intermediate NO release, for example organic nitrates. The drawback of this treatment is the development of tolerance and a reduction of activity, and the higher dosage which is required because of this.

10 Various sGC stimulators which do not act via NO release were described by Vesely in a series of publications. However, the compounds, most of which are hormones, plant hormones, vitamins or natural compounds such as, for example, lizard poisons predominantly only have weak effects on the cGMP formation in cell lysates. D. L. Vesely, *Eur. J. Clin. Invest.*, vol. 15, 1985, p. 258; D. L. Vesely, *Biochem. Biophys. Res. Comm.*, vol. 88, 1979, p. 1244. A stimulation of heme-free guanylate cyclase by  
15 protoporphyrin IX was demonstrated by Ignarro et al., *Adv. Pharmacol.*, vol. 26, 1994, p. 35. Pettibone et al., *Eur. J. Pharmacol.*, vol. 116, 1985 p. 307, described an antihypertensive action of diphenyliodonium hexafluorophosphate and attributed this to a stimulation of sGC. According to Yu et al., *Brit. J. Pharmacol.*, vol. 114, 1995, p. 1587,  
20 isoliquiritigenin, which has a relaxing action on isolated rat aortas, also activates sGC. Ko et al., *Blood* vol. 84, 1994, p. 4226, Yu et al., *Biochem. J.* vol. 306, 1995, p. 787, and


Wu et al., *Brit. J. Pharmacol.* vol. 116, 1995, p. 1973, demonstrated a sGC-stimulating activity of 1-benzyl-3-(5-hydroxymethyl-2-furyl)indazole and demonstrated an antiproliferative and thrombocyte-inhibiting action. Pyrazoles and fused pyrazoles which exhibit a sGC-stimulating activity are described in European Patent Application No. 5 908,456 and German Patent Application No. 19,744,027.

A series of 2-sulfonylaminobenzoic acid N-arylamides, the N-aryl group of which carries a thio substituent, have been mentioned in the literature. These compounds in which the N-aryl group generally carries as further substituents groups which are readily oxidizable  
10 such as, for example, two hydroxy groups being in para position with respect to one another and which in this case can be regarded as hydroquinone derivatives, are auxiliaries for the preparation of photographic materials (see, for example, Chemical Abstracts 119, 105757; 120, 41858; 123, 70224; or 126, 257007). If isolated structural elements are considered then the N-aryl group in these known compounds corresponds  
15 to the group  $R^1-S(O)_n-A^1$  in formula I in case  $A^1$  denotes a 1,4-phenylene residue which in positions 2 and 5 carries hydroxy groups (or oxy substituents), and the number  $n$  is 0. British patent publication No. 876,526 (Chemical Abstracts 56, 15432e) discloses 3,5-dichloro-2-methylsulfonylaminobenzoic acid N-(5-chloro-2-(4-chlorophenylmercapto)-phenyl)-amide which can be used for the protection of wool against moths. Compounds  
20 covered by British patent publication No. 876,526 correspond to compounds of the formula I if simultaneously the ring  $A^1$  which comprises the carbon atoms which carry

the groups  $C(=X)-NH-$  and  $NH-SO_2R^2$ , together with the residues  $R^3$ , is a benzene ring which carries one to four halogen atoms from the series chlorine and bromine,  $R^2$  is  $(C_1-C_4)$ -alkyl,  $X$  is oxygen and the group  $R^1-S(O)_n-A^1-$  is a phenylmercaptophenyl- residue (= phenylthiophenyl-) which is substituted by halogen and/or trifluoromethyl and which can  
5 also be substituted by methyl or  $(C_1-C_4)$ -alkoxy, and the total number of halogen atoms and trifluoromethyl groups is greater than two. Pharmacological activities of these known 2-sulfonylaminobenzoic acid N-arylamides are not disclosed.

Surprisingly, it has now been found that the compounds of the present invention effect a  
10 strong activation of guanylate cyclase and are therefore suitable for the therapy and prophylaxis of disorders which are associated with a low cGMP level.

Thus, the present invention relates to compounds of the formula I:



wherein

A¹ is a divalent residue from the series phenylene, naphthylene and heteroarylene which can all be substituted by one or more identical or different substituents from the series halogen, (C₁-C₅)-alkyl, phenyl, tolyl, CF₃, NO₂, OH, -O-(C₁-C₅)-alkyl, -O-(C₂-C₄)-alkyl-O-(C₁-C₃)-alkyl, (C₁-C₂)-alkylenedioxy, NH₂, -NH-(C₁-C₃)-alkyl, -N((C₁-C₃)-alkyl)₂, -NH-CHO, -NH-CO-(C₁-C₅)-alkyl, -CN, -CO-NH₂, -CO-NH-(C₁-C₃)-alkyl, -CO-N((C₁-C₃)-alkyl)₂, -CO-OH, -CO-O-(C₁-C₅)-alkyl, heterocyclyl, CHO, -CO-(C₁-C₅)-alkyl, -S(O)ₙ-(C₁-C₄)-alkyl, -S(O)ₙ-phenyl and -S(O)ₙ-tolyl;

the ring A² which comprises the carbon atoms which carry the groups C(=X)-NH- and

NH-SO₂R² is a benzene ring, a naphthalene ring, a saturated or partially unsaturated 3-membered to 7-membered carbocycle, a saturated or partially unsaturated or aromatic



monocyclic 5-membered to 7-membered heterocycle which contains one or more ring heteroatoms from the series N, O and S, or a saturated or partially unsaturated or aromatic bicyclic 8-membered to 10-membered heterocycle which contains one or more ring heteroatoms from the series N, O and S;

5

$R^1$  is aryl, heterocyclyl or  $(C_1-C_{18})$ -alkyl which can be substituted by one or more identical or different residues  $R^4$  or, if the number  $n$  in the group  $R^1-S(O)_n-$  is 2,  $R^1$  can also be  $NR^5R^6$  or, if the number  $n$  in the group  $R^1-S(O)_n-$  is 0,  $R^1$  can also be  $-CN$ ;

10  $R^2$  is aryl, heterocyclyl,  $NR^5R^6$  or  $(C_1-C_{10})$ -alkyl which can be substituted by one or more identical or different residues  $R^4$ ;

$R^3$  denotes one or more identical or different residues from the series hydrogen, halogen,  $CF_3$ ,  $OH$ ,  $-O-(C_1-C_7)$ -alkyl,  $-O-(C_2-C_4)$ -alkyl- $O-(C_1-C_7)$ -alkyl,  $-O$ -aryl,  $(C_1-C_2)$ -alkylenedioxy,  $NO_2$ ,  $-CN$ ,  $NR^7R^8$ ,  $-CO-NR^7R^8$ ,  $-CO-OH$ ,  $-CO-O-(C_1-C_5)$ -alkyl, 15  $heterocyclyl$ ,  $-S(O)_n-(C_1-C_5)$ -alkyl and  $(C_1-C_5)$ -alkyl which can be substituted by one or more identical or different residues  $R^4$ ;

$R^4$  is fluorine,  $OH$ ,  $-O-(C_1-C_{10})$ -alkyl,  $-O-(C_2-C_4)$ -alkyl- $O-(C_1-C_7)$ -alkyl,  $-O$ -aryl,  $-CN$ , 20  $NR^7R^8$ ,  $-CO-NH_2$ ,  $-CO-NH-(C_1-C_3)$ -alkyl,  $-CO-N((C_1-C_3)$ -alkyl) $_2$ ,  $-CO-OH$ ,  $-CO-O-(C_1-C_5)$ -alkyl, heterocyclyl or oxo;

$R^5$  is hydrogen,  $(C_1-C_{10})$ -alkyl which can be substituted by one or more identical or different substituents  $R^4$  and/or by aryl, or is aryl, heterocyclyl,  $-CO-NR^7R^8$ ,  $-CO$ -aryl or  $-CO-(C_1-C_{10})$ -alkyl wherein the alkyl residue can be substituted by one or more identical or different residues  $R^4$ ;

5

$R^6$  independently of  $R^5$  has one of the meanings indicated for  $R^5$ , or

$R^5$  and  $R^6$  together with the nitrogen atom to which they are bonded form a 5-membered to 8-membered saturated or partially unsaturated ring which in addition to the nitrogen atom which carries the groups  $R^5$  and  $R^6$  can contain one or more further ring

10 heteroatoms from the series N, O and S and which can be substituted by one or more identical or different substituents from the series fluorine,  $(C_1-C_5)$ -alkyl, hydroxy- $(C_1-C_3)$ -alkyl-,  $-(C_1-C_3)$ -alkyl-O- $(C_1-C_4)$ -alkyl, aryl,  $CF_3$ , OH,  $-O-(C_1-C_7)$ -alkyl,  $-O$ -aryl,  $-O-(C_2-C_4)$ -alkyl-O- $(C_1-C_7)$ -alkyl,  $(C_2-C_3)$ -alkylenedioxy,  $NR^7R^8$ ,  $-CN$ ,  $-CO-NH_2$ ,  $-CO-NH-(C_1-C_3)$ -alkyl,  $-CO-N((C_1-C_3)$ -alkyl) $_2$ ,  $-CO-OH$ ,  $-CO-O-(C_1-C_5)$ -alkyl, CHO,  $-CO-(C_1-C_5)$ -alkyl,

15  $-S(O)_n-(C_1-C_4)$ -alkyl,  $-S(O)_n-NH_2$ ,  $-S(O)_n-NH-(C_1-C_3)$ -alkyl,  $-S(O)_n-N((C_1-C_3)$ -alkyl) $_2$ , oxo,  $-(CH_2)_m-NH_2$ ,  $-(CH_2)_m-NH-(C_1-C_4)$ -alkyl and  $-(CH_2)_m-N((C_1-C_4)$ -alkyl) $_2$  where in the substituent  $-(CH_2)_m-N((C_1-C_4)$ -alkyl) $_2$  the two alkyl groups can be connected by a single bond and then together with the nitrogen atom carrying them form a 5-membered to 7-membered ring which besides that nitrogen atom and the carbon atoms can additionally

20 contain an oxygen atom, a sulfur atom or a group  $NR^5$  as ring member;

R<sup>7</sup> is hydrogen or (C<sub>1</sub>-C<sub>7</sub>)-alkyl which can be substituted by one or more identical or different substituents from the series OH, -O-(C<sub>1</sub>-C<sub>5</sub>)-alkyl, NH<sub>2</sub>, -NH-(C<sub>1</sub>-C<sub>4</sub>)-alkyl and -N((C<sub>1</sub>-C<sub>4</sub>)-alkyl)<sub>2</sub> where in the substituent N((C<sub>1</sub>-C<sub>4</sub>)-alkyl)<sub>2</sub> the two alkyl groups can be connected by a single bond and then together with the nitrogen atom carrying them

5 form a 5-membered to 7-membered ring which besides that nitrogen atom and the carbon atoms can additionally contain an oxygen atom, a sulfur atom or a group NR<sup>5</sup> as ring member;

R<sup>8</sup> independently of R<sup>7</sup> has one of the meanings of R<sup>7</sup> or is -CO-(C<sub>1</sub>-C<sub>4</sub>)-alkyl;

10 "aryl" is phenyl, naphthyl or heteroaryl which can all be substituted by one or more identical or different substituent from the series halogen, (C<sub>1</sub>-C<sub>5</sub>)-alkyl, phenyl, tolyl, CF<sub>3</sub>, -O-CF<sub>3</sub>, NO<sub>2</sub>, OH, -O-(C<sub>1</sub>-C<sub>5</sub>)-alkyl, -O-(C<sub>2</sub>-C<sub>4</sub>)-alkyl-O-(C<sub>1</sub>-C<sub>3</sub>)-alkyl, (C<sub>1</sub>-C<sub>2</sub>)-alkylenedioxy, NH<sub>2</sub>, -NH-(C<sub>1</sub>-C<sub>3</sub>)-alkyl, -N((C<sub>1</sub>-C<sub>3</sub>)-alkyl)<sub>2</sub>, -NH-CHO, -NH-CO-(C<sub>1</sub>-C<sub>5</sub>)-alkyl, -CN, -CO-NH<sub>2</sub>, -CO-NH-(C<sub>1</sub>-C<sub>3</sub>)-alkyl, -CO-N((C<sub>1</sub>-C<sub>3</sub>)-alkyl)<sub>2</sub>, -CO-OH, -CO-O-(C<sub>1</sub>-C<sub>5</sub>)-alkyl, heterocyclyl, CHO, -CO-(C<sub>1</sub>-C<sub>5</sub>)-alkyl, -S(O)<sub>n</sub>-(C<sub>1</sub>-C<sub>4</sub>)-alkyl, -S(O)<sub>n</sub>-phenyl and -S(O)<sub>n</sub>-tolyl;

20 "heteroaryl" and "heteroarylene" are a residue of a monocyclic 5-membered or 6-membered aromatic heterocycle or of a bicyclic 8-membered to 10-membered aromatic

heterocycle each of which contains one or more ring heteroatoms from the series N, O and S;

"heterocyclyl" is a residue of a monocyclic or polycyclic 5-membered to 11-membered saturated or partially unsaturated heterocycle which contains one or more ring heteroatoms from the series N, O and S and which can be substituted by one or more identical or different substituents from the series fluorine, (C<sub>1</sub>-C<sub>5</sub>)-alkyl, OH, -O-(C<sub>1</sub>-C<sub>5</sub>)-alkyl, -O-(C<sub>2</sub>-C<sub>4</sub>)-alkyl-O-(C<sub>1</sub>-C<sub>3</sub>)-alkyl, NH<sub>2</sub>, -NH-(C<sub>1</sub>-C<sub>3</sub>)-alkyl, -N((C<sub>1</sub>-C<sub>3</sub>)-alkyl)<sub>2</sub>, -CN, -CO-NH<sub>2</sub>, -CO-NH-(C<sub>1</sub>-C<sub>3</sub>)-alkyl, -CO-N((C<sub>1</sub>-C<sub>3</sub>)-alkyl)<sub>2</sub>, -CO-OH and -CO-O-(C<sub>1</sub>-C<sub>5</sub>)-alkyl;

n is 0, 1 or 2;

m is 2, 3 or 4;

X is O or NH or X is a nitrogen atom which via a single bond is attached to a ring carbon atom in the group A<sup>1</sup> which ring carbon atom is directly adjacent to the carbon atom in A<sup>1</sup> carrying the group -NH-C(=X)- so that the group -NH-C(=X)- together with the carbon atoms in A<sup>1</sup> carrying it forms an anellated imidazole ring;

in all their stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts;

where, however, the compound of formula I is excluded wherein simultaneously the ring  $A^2$  which comprises the carbon atoms which carry the groups  $C(=X)-NH-$  and  $NH-SO_2R^2$  is a benzene ring which is substituted in positions 3 and 5 by chlorine,  $R^2$  is methyl, X is oxygen and  $R^1-S(O)_n-A^1-$  is a 5-chloro-2-(4-chlorophenylmercapto)-phenyl residue.

5

If groups or substituents can occur several times in the compounds of formula I such as, for example  $R^3$ ,  $R^4$ ,  $R^5$ , aryl, heterocyclyl, alkyl, or the numbers n and m, they can all independently of one another have the meanings indicated and can in each case be identical or different.

10

Alkyl residues can be straight-chain or branched. This also applies when they are part of other groups, for example in alkoxy groups, alkoxycarbonyl groups or amino groups, or when they are substituted. Examples of alkyl groups are methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl,

15 pentadecyl, hexadecyl, heptadecyl, octadecyl, the n-isomers of these residues, isopropyl, isobutyl, isopentyl, sec-butyl, tert-butyl, neopentyl, 3,3-dimethylbutyl. The term alkyl here also expressly includes unsaturated alkyl residues, i. e. alkyl residues which contain one or more double bonds and/or one or more triple bonds such as, for example, alkenyl residues and alkynyl residues. Of course, an unsaturated alkyl group  
20 contains at least two carbon atoms. Specific alkyl groups whose number of carbon atoms can vary from 1 to a given upper limit, thus also comprise unsaturated alkyl

groups whose number of carbon atoms can vary from 2 to the given upper limit.

Examples of such residues are the vinyl residue, the 1-propenyl residue, the 2-propenyl residue (allyl residue), the 2-butenyl residue, the 2-methyl-2-propenyl residue, the 3-methyl-2-butenyl residue, the ethynyl residue, the 2-propinyl residue (propargyl residue),  
5 the 2-butylnyl residue or the 3-butylnyl residue. Further, the term alkyl here also expressly includes alkyl residues in which by an internal ring closure within the alkyl group a cyclic system is formed, i. e. the term alkyl also includes saturated and partially unsaturated cycloalkyl residues and cycloalkyl-alkyl- residues (alkyl substituted by cycloalkyl). Of course, a monocyclic cycloalkyl group contains at least three carbon atoms.

Specific alkyl groups whose number of carbon atoms can vary from 1 to a given upper limit, thus also comprise monocyclic cycloalkyl groups whose number of carbon atoms can vary from 3 to the given upper limit, and appropriate cycloalkyl-alkyl- groups.

Examples of such cycloalkyl residues are cyclopropyl, cyclobutyl, cyclopentyl,

15 cyclohexyl, cycloheptyl and cyclooctyl which can all also be substituted by one or more identical or different ( $C_1$ - $C_4$ )-alkyl residues, in particular by methyl. Examples of such substituted cycloalkyl residues are 4-methylcyclohexyl, 4-tert-butylcyclohexyl or 2,3-dimethylcyclopentyl. Furthermore, unless stated otherwise the term alkyl here also expressly includes unsubstituted alkyl residues as well as alkyl residues which are  
20 substituted by one or more, for example one, two, three or four, identical or different aryl residues. The term alkyl thus here also expressly includes arylalkyl- residues such as,

for example, aryl-(C<sub>1</sub>-C<sub>4</sub>)-alkyl-, for example benzyl residues, phenylethyl residues or indanyl residues. In substituted alkyl residues, for example arylalkyl-, hydroxyalkyl- such as -(C<sub>1</sub>-C<sub>3</sub>)-alkyl-OH or alkoxyalkyl- such as -(C<sub>1</sub>-C<sub>3</sub>)-alkyl-O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl, the substituents can be present in any desired position.

5

A saturated or partially unsaturated 3-membered to 7-membered carbocycle representing the ring A<sup>2</sup> can be derived from the monocyclic parent systems cyclopropane, cyclobutane, cyclopentane, cyclohexane or cycloheptane. If the carbocycle is unsaturated it can contain, for example, one double bond or, in the case of a 5-membered ring, 6-membered ring or 7-membered ring, also two double bonds which can be isolated or conjugated. Double bonds can be present in any positions with respect to the groups C(=X)-NH- and NH-SO<sub>2</sub>-R<sup>2</sup>, i.e., for example a double bond can also be present between the two ring carbon atoms which carry these two groups.

15 Unless stated otherwise, phenyl residues, naphthyl residues and heterocyclic residues, for example heteroaryl residues, can be unsubstituted or can carry one or more, for example one, two, three or four, identical or different substituents which can be in any desired positions. Unless stated otherwise, in these residues for example those substituents can be present which are indicated as substituents of an aryl group. A  
20 preferred series of substituents that can be present in the residue aryl is formed by the substituents halogen, (C<sub>1</sub>-C<sub>5</sub>)-alkyl, phenyl, tolyl, CF<sub>3</sub>, NO<sub>2</sub>, OH, -O-(C<sub>1</sub>-C<sub>5</sub>)-alkyl, -O-(C<sub>2</sub>-

$C_4$ )-alkyl-O-( $C_1$ - $C_3$ )-alkyl, ( $C_1$ - $C_2$ )-alkylenedioxy,  $NH_2$ , -NH-( $C_1$ - $C_3$ )-alkyl, -N(( $C_1$ - $C_3$ )-alkyl)<sub>2</sub>, -NH-CHO, -NH-CO-( $C_1$ - $C_5$ )-alkyl, -CN, -CO-NH<sub>2</sub>, -CO-NH-( $C_1$ - $C_3$ )-alkyl, -CO-N(( $C_1$ - $C_3$ )-alkyl)<sub>2</sub>, -CO-OH, -CO-O-( $C_1$ - $C_5$ )-alkyl, heterocyclyl, CHO, -CO-( $C_1$ - $C_5$ )-alkyl, -S(O)<sub>n</sub>-( $C_1$ - $C_4$ )-alkyl, -S(O)<sub>n</sub>-phenyl and -S(O)<sub>n</sub>-tolyl. If in compounds of the formula

5 I nitro groups are present as substituents, in total only up to two nitro groups can be present in the molecules. If phenyl residues, phenoxy residues, benzyl residues or benzyloxy residues are present as substituents in, for example, aryl residues like phenyl residues and/or in heterocyclic residues then in these substituents the benzene ring can also be unsubstituted or substituted by one or more, for example one, two, three or four, identical or different residues, for example by residues from the series ( $C_1$ - $C_4$ )-alkyl, 10 halogen, hydroxy, ( $C_1$ - $C_4$ )-alkoxy, trifluoromethyl, cyano, hydroxycarbonyl, (( $C_1$ - $C_4$ )-alkoxy)carbonyl, aminocarbonyl, nitro, amino, ( $C_1$ - $C_4$ )-alkylamino, di-(( $C_1$ - $C_4$ )-alkyl)amino and (( $C_1$ - $C_4$ )-alkyl)carbonylamino.

15 In monosubstituted phenyl residues the substituent can be in the 2-position, the 3-position or the 4-position, in disubstituted phenyl residues the substituents can be in 2,3-position, 2,4-position, 2,5-position, 2,6-position, 3,4-position or 3,5-position. In trisubstituted phenyl residues the substituents can be in 2,3,4-position, 2,3,5-position, 2,3,6-position, 2,4,5-position, 2,4,6-position or 3,4,5-position. Toly (i.e., methylphenyl) 20 can be 2-tolyl, 3-tolyl or 4-tolyl. Naphthyl can be 1-naphthyl or 2-naphthyl. In monosubstituted 1-naphthyl residues the substituent can be in the 2-position, the 3-



position, the 4-position, the 5-position, the 6-position, the 7-position or the 8-position, in monosubstituted 2-naphthyl residues in the 1-position, the 3-position, the 4-position, the 5-position, the 6-position, the 7-position or the 8-position.

- 5 The above explanations as well as the following explanations relating to monovalent residues correspondingly apply to the divalent residues phenylene, naphthylene and heteroarylene. The free bonds via which the divalent residues are attached to the adjacent groups can be present on any ring carbon atoms. In the case of a phenylene residue they can be in 1,2-position (ortho-phenylene), 1,3-position (meta-phenylene) or 1,4-position (para-phenylene). In the case of a naphthylene residue the free bonds can be in 1,2-position (= 1,2-naphthylene or 1,2-naphthalinediyl) or in 1,3-position, 1,4-position, 1,5-position, 1,6-position, 1,7-position, 1,8-position, 2,3-position, 2,6-position or 2,7-position. In the case of 5-membered ring aromatics containing one heteroatom such as, for example, thiophene or furan, the two free bonds can be in 2,3-position, 2,4-position, 2,5-position or 3,4-position. A divalent residue derived from pyridine can be a 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-pyridinediyl residue. In the case of unsymmetrical divalent residues the present invention includes all positional isomers, i.e., in the case of a 2,3-pyridinediyl residue, for example, it includes the compound in which the one adjacent group is present in the 2-position and the other adjacent group is present in the 3-position as well as the compound in which the one adjacent group is present in the 3-position and the other adjacent group is present in the 2-position.

Heteroaryl residues, heteroarylene residues, heterocyclyl residues, heterocycles representing the ring A<sup>2</sup> and rings which are formed by two groups bonded to a nitrogen atom together with this nitrogen atom are preferably derived from heterocycles which contain one, two, three or four identical or different ring heteroatoms, more preferably from heterocycles which contain one, two or three, in particular one or two, identical or different heteroatoms. Unless stated otherwise, the heterocycles can be monocyclic or polycyclic, for example monocyclic, bicyclic or tricyclic. Preferably they are monocyclic or bicyclic. The rings preferably are 5-membered rings, 6-membered ring or 7-membered ring. Examples of monocyclic and bicyclic heterocyclic systems from which residues occurring in the compounds of the formula I can be derived, are pyrrole, furan, thiophene, imidazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, 1,3-dioxole, 1,3-oxazole (= oxazole), 1,2-oxazole (= isoxazole), 1,3-thiazole (= thiazole), 1,2-thiazole (= isothiazole), tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, pyran, thiopyran, 1,4-dioxine, 1,2-oxazine, 1,3-oxazine, 1,4-oxazine, 1,2-thiazine, 1,3-thiazine, 1,4-thiazine, 1,2,3-triazine, 1,2,4-triazine, 1,3,5-triazine, 1,2,4,5-tetrazine, azepine, 1,2-diazepine, 1,3-diazepine, 1,4-diazepine, 1,3-oxazepine, 1,3-thiazepine, indole, benzothiophene, benzofuran, benzothiazole, benzimidazole, quinoline, isoquinoline, cinnoline, quinazoline, quinoxaline, phthalazine, thienothiophenes, 1,8-naphthyridine and other naphthyridines, pteridin, or phenothiazine, each of them in saturated form (perhydro form) or in partially unsaturated form (for example in the dihydro form or the tetrahydro form) or in maximally unsaturated form, insofar as the respective forms are known and stable.

Thus, the heterocycles which are suitable also include, for example, the saturated heterocycles pyrrolidine, piperidine, piperazine, morpholine and thiomorpholine. The degree of saturation of heterocyclic groups is indicated in their individual definitions. Unsaturated heterocycles can contain, for example, one, two or three double bonds within the ring system. 5-membered rings and 6-membered rings can in particular also be aromatic.

The residues derived from these heterocycles can be attached via any suitable carbon atom. Nitrogen heterocycles which can carry a hydrogen atom or a substituent on a ring nitrogen atom, for example pyrrole, imidazole, pyrrolidine, morpholine, piperazine etc., can also be attached via a ring nitrogen atom, in particular if the heterocyclic residue in question is bonded to a carbon atom. For example, a thienyl residue can be present as 2-thienyl residue or 3-thienyl residue, a furyl residue as 2-furyl residue or 3-furyl residue, a pyridyl residue as 2-pyridyl residue, 3-pyridyl residue or 4-pyridyl residue, a piperidinyl residue as 1-piperidinyl residue (i.e., piperidino residue), 2-piperidinyl residue, 3-piperidinyl residue or 4-piperidinyl residue, a (thio)morpholinyl residue as 2-(thio)morpholinyl residue, 3-(thio)morpholinyl residue or 4-(thio)morpholinyl residue (i.e., thiomorpholino residue). A residue derived from 1,3-thiazole or imidazole which is attached via a carbon atom can be attached via the 2-position, the 4-position or the 5-position.

Unless stated otherwise the heterocyclic groups can be unsubstituted or can carry one or more, for example one, two, three, or four, identical or different substituents.

Substituents in heterocycles can be present in any desired positions, for example in a 2-thienyl residue or 2-furyl residue in the 3-position and/or in the 4-position and/or in the 5-

position, in a 3-thienyl residue or 3-furyl residue in the 2-position and/or in the 4-position and/or in the 5-position, in a 2-pyridyl residue in the 3-position and/or in the 4-position

and/or in the 5-position and/or in the 6-position, in a 3-pyridyl residue in the 2-position and/or in the 4-position and/or in the 5-position and/or in the 6-position, in a 4-pyridyl

residue in the 2-position and/or in the 3-position and/or in the 5-position and/or in the 6-position. Unless stated otherwise, for example those substituents can be present as

substituents in heterocyclic groups which are indicated in the definition of the group aryl, and in the case of saturated and partially unsaturated heterocycles as further

substituents also the oxo group and the thioxo group can be present. Substituents on a heterocycle as well as substituents on a carbocycle can also form a ring, i.e., to a ring

system further rings can be condensed (or anellated) so that, for example, also

cyclopenta-condensed, cyclohexa-condensed or benzo-condensed rings can be

present. Suitable substituents on a substitutable ring nitrogen atom of a heterocycle are

in particular, for example, unsubstituted (C<sub>1</sub>-C<sub>5</sub>)-alkyl residues and aryl-substituted alkyl residues, aryl residues, acyl residues such as -CO-(C<sub>1</sub>-C<sub>5</sub>)-alkyl, or sulfonyl residues

such as -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>5</sub>)-alkyl. Suitable nitrogen heterocycles can also be present as N-



oxides or as quaternary salts containing a counterion which is derived from a





physiologically acceptable acid. Pyridyl residues, for example, can be present as pyridine-N-oxides.



"Halogen" is fluorine, chlorine, bromine or iodine, and preferably fluorine or chlorine.

5

Without limiting the present invention, in the formulae Ia, Ib, Ic, Id, Ie, If, Ig and Ih examples of groups of compounds of the invention are shown in which A<sup>2</sup> in the formula I has specific denotations. A<sup>1</sup>, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, X and n in the formulae Ia, Ib, Ic, Id, Ie, If, Ig and Ih are defined as above for the formula I, and the number k in the formula Ib is 1, 2, 3, 4 or 5, in particular 3 or 4.








On the benzene ring depicted in formula Ia which carries the groups C(=X)-NH- and -NHSO<sub>2</sub>R<sup>2</sup>, four positions are present which can carry a residue R<sup>3</sup>. The compounds of formula Ia can thus carry four residues R<sup>3</sup> which, independently of one another, can all be hydrogen or can have a meaning different from hydrogen, i.e., in the compounds of formula Ia the benzene ring depicted in formula Ia can be unsubstituted or can carry one, two, three or four identical or different substituents from the series halogen, CF<sub>3</sub>, OH, -O-(C<sub>1</sub>-C<sub>7</sub>)-alkyl, -O-(C<sub>2</sub>-C<sub>4</sub>)-alkyl-O-(C<sub>1</sub>-C<sub>7</sub>)-alkyl, -O-aryl, (C<sub>1</sub>-C<sub>2</sub>)-alkylenedioxy, NO<sub>2</sub>, -CN, NR<sup>7</sup>R<sup>8</sup>, -CO-NR<sup>7</sup>R<sup>8</sup>, -CO-OH, -CO-O-(C<sub>1</sub>-C<sub>5</sub>)-alkyl, heterocyclyl, -S(O)<sub>n</sub>-(C<sub>1</sub>-C<sub>5</sub>)-alkyl and (C<sub>1</sub>-C<sub>5</sub>)-alkyl which can be substituted by one or more identical or different residues R<sup>4</sup>. These explanations accordingly also apply to the compounds of formulae Ib to Ih.

The compounds of the formula I wherein X is a nitrogen atom which via a single bond is attached to a ring carbon atom in the group A<sup>1</sup> which ring carbon atom is directly adjacent to the carbon atom in A<sup>1</sup> carrying the group -NH-C(=X)- so that the group -NH-C(=X)- together with the carbon atoms in A<sup>1</sup> carrying it forms an anellated

5 imidazole ring, are represented by the formula li.



15 A<sup>2</sup>, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and n in the formula li are defined as above for the formula I. The ring A<sup>3</sup> which has resulted from the group A<sup>1</sup> by the formation of a bond to the nitrogen atom representing X and which ring comprises the two carbon atoms depicted in formula li carrying the nitrogen atoms of the anellated imidazole ring, is a benzene ring, a naphthalene ring or a heteroaromatic ring, where to these rings the above explanations

20 relating to A<sup>1</sup> correspondingly apply.



The present invention includes all stereoisomeric forms of the compounds of the formula

I. Centers of asymmetry that are present in the compounds of formula I can all independently of one another have S configuration or R configuration. The invention includes all possible enantiomers and diastereomers and mixtures of two or more

5 stereoisomers, for example mixtures of enantiomers and/or diastereomers, in all ratios.

Thus, enantiomers are a subject of the invention in enantiomerically pure form, both as levorotatory and as dextrorotatory antipodes, in the form of racemates and in the form of

mixtures of the two enantiomers in all ratios. In the case of a cis/trans isomerism the invention includes both the cis form and the trans form as well as mixtures of these

10 forms in all ratios. The preparation of individual stereoisomers can be carried out, if desired, by separation of a mixture by customary methods, for example by

chromatography or crystallization, by the use of stereochemically uniform starting

materials for the synthesis or by stereoselective synthesis. Optionally a derivatization

can be carried out before a separation of stereoisomers. The separation of a mixture of

15 stereoisomers can be carried out at the stage of the compounds of the formula I or at

the stage of an intermediate during the synthesis. The present invention also includes

all tautomeric forms of the compounds of formula I.

If the compounds of the formula I contain one or more acidic or basic groups the

20 invention also includes the corresponding physiologically or toxicologically acceptable

salts, in particular the pharmaceutically utilizable salts. Thus, the compounds of the

formula I which contain acidic groups can be present on these groups and can be used according to the invention, for example, as alkali metal salts, alkaline earth metal salts or as ammonium salts. Examples of such salts are sodium salts, potassium salts, calcium salts, magnesium salts or salts with ammonia or organic amines such as, for example, ethylamine, ethanolamine, triethanolamine or amino acids. Compounds of the formula I which contain one or more basic groups, i.e. groups which can be protonated, can be present and can be used according to the invention in the form of their acid addition salts with inorganic or organic acids, for example as salts with hydrogen chloride, hydrogen bromide, phosphoric acid, sulfuric acid, nitric acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acids, oxalic acid, acetic acid, tartaric acid, lactic acid, salicylic acid, benzoic acid, formic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, malic acid, sulfaminic acid, phenylpropionic acid, gluconic acid, ascorbic acid, isonicotinic acid, citric acid, adipic acid, etc. If the compounds of the formula I simultaneously contain acidic and basic groups in the molecule the invention also includes, in addition to the salt forms mentioned, inner salts or betaines (zwitterions). Salts can be obtained from the compounds of the formula I by customary methods which are known to the person skilled in the art, for example by combination with an organic or inorganic acid or base in a solvent or dispersant, or by anion exchange or cation exchange from other salts. The present invention also includes all salts of the compounds of the formula I which, owing to low physiological compatibility, are not

directly suitable for use in pharmaceuticals but which can be used, for example, as intermediates for chemical reactions or for the preparation of physiologically acceptable salts.

- 5 The present invention furthermore includes all solvates of compounds of the formula I, for example hydrates or adducts with alcohols, and also derivatives of the compounds of the formula I such as, for example esters, amides, prodrugs and active metabolites.

A<sup>1</sup> preferably is a phenylene residue or a 5-membered or 6-membered heteroarylene residue, more preferably phenylene, where all these residues can be unsubstituted or can be substituted as indicated. If the group A<sup>1</sup> is substituted, i. e. if it carries one or more further substituents in addition to the group R<sup>1</sup>-S(O)<sub>n</sub>, it is preferably substituted by one or two identical or different of the substituents indicated above. Preferably a phenylene residue representing A<sup>1</sup> is unsubstituted, i. e. besides the groups R<sup>1</sup>-S(O)<sub>n</sub> and C(=X)-NH it carries four hydrogen atoms. The group R<sup>1</sup>-S(O)<sub>n</sub> is preferably attached to a carbon atom in A<sup>1</sup> which is not directly adjacent to the carbon atom which carries the group C(=X)-NH. If A<sup>1</sup> is phenylene the group R<sup>1</sup>-S(O)<sub>n</sub> is particularly preferably located in the meta position or in the para position, more particularly preferably in the para position, with respect to the carbon atom which carries the group C(=X)-NH.

The ring A<sup>2</sup> which comprises the two carbon atoms which carry the groups R<sup>2</sup>-SO<sub>2</sub>-NH and C(=X)-NH- preferably is an aromatic ring, more preferably a benzene ring or a thiophene ring, particularly preferably a benzene ring, where all these rings can be unsubstituted or substituted by one or more residues R<sup>3</sup> which are different from  
5 hydrogen.

R<sup>1</sup> preferably is (C<sub>1</sub>-C<sub>7</sub>)-alkyl, NR<sup>5</sup>R<sup>6</sup> or aryl, more preferably NR<sup>5</sup>R<sup>6</sup>, phenyl or 5-membered or 6-membered heteroaryl, particularly preferably NR<sup>5</sup>R<sup>6</sup>, where all these residues can be unsubstituted or substituted as indicated and where, as stated above, R<sup>1</sup> can be NR<sup>5</sup>R<sup>6</sup> if the number n in the group R<sup>1</sup>-S(O)<sub>n</sub>- is 2.

R<sup>2</sup> preferably is aryl, more preferably phenyl or 5-membered or 6-membered heteroaryl, particularly preferably phenyl or a residue of a monocyclic 5-membered or 6-membered aromatic heterocycle which contains one or two identical or different heteroatoms from  
15 the series N, O and S such as, for example, phenyl, thienyl, pyrazolyl, imidazolyl, isoxazolyl, thiazolyl, pyridyl etc., in particular phenyl or 2-thienyl, where all these residues can be unsubstituted or substituted as indicated. Preferably an aryl residue representing R<sup>2</sup> is substituted. If an aryl residue representing R<sup>2</sup> is substituted it is preferably substituted by one, two or three, in particular by one or two, identical or  
20 different substituents. Substituents in an aryl residue representing R<sup>2</sup> preferably are substituents from the series halogen, CF<sub>3</sub>, -O-CF<sub>3</sub>, NO<sub>2</sub>, -CN, (C<sub>1</sub>-C<sub>4</sub>)-alkyl and -O-(C<sub>1</sub>-

C<sub>4</sub>)-alkyl, more preferably substituents from the series F, Cl, Br, CF<sub>3</sub>, -O-CF<sub>3</sub>, NO<sub>2</sub>, -CN, CH<sub>3</sub> and -OCH<sub>3</sub>. Especially preferably a substituted aryl residue representing R<sup>2</sup> is substituted by Cl, for example by one or two, in particular one, chlorine atoms.

- 5 The rings representing A<sup>2</sup> can be unsubstituted or substituted as indicated. When they are unsubstituted they only carry residues R<sup>3</sup> which are hydrogen. When they are substituted they carry one or more residues R<sup>3</sup> which are different from hydrogen. Those substituent positions which do not carry a residue R<sup>3</sup> which is different from hydrogen, carry hydrogen atoms. If the ring A<sup>2</sup> carries one or more residues R<sup>3</sup> which are different from hydrogen it preferably carries one or two such residues R<sup>3</sup>, in particular one such residue R<sup>3</sup>. Residues R<sup>3</sup> which are different from hydrogen are preferably located in positions of the ring A<sup>2</sup> which are not directly adjacent to the groups C(=X)-NH and R<sup>2</sup>-SO<sub>2</sub>-NH. If A<sup>2</sup> is a saturated or partially unsaturated carbocycle, residues R<sup>3</sup> which are different from hydrogen preferably are (C<sub>1</sub>-C<sub>4</sub>)-alkyl, in particular methyl. If A<sup>2</sup> is an aromatic ring, in particular if A<sup>2</sup> is a benzene ring, residues R<sup>3</sup> which are different from hydrogen preferably are (C<sub>1</sub>-C<sub>3</sub>)-alkyl, halogen, (C<sub>1</sub>-C<sub>3</sub>)-alkoxy or CF<sub>3</sub>, more preferably methyl, chlorine or methoxy. If A<sup>2</sup> is an aromatic ring, in particular a benzene ring, it is very particularly preferred if the ring carries one chlorine atom or two methoxy groups as substituents, i. e., if one residue R<sup>3</sup> is present which is chlorine or if two residues R<sup>3</sup> are present which are methoxy, and the other positions on the benzene ring carry hydrogen atoms. If A<sup>2</sup> is a benzene ring residues R<sup>3</sup>

which are different from hydrogen are preferably located in positions 4 and/or 5 (with respect to the group  $C(=X)-NH$  in the 1-position and the group  $R^2-SO_2-NH$  in the 2-position).

5 If a group is substituted by one or more residues  $R^4$  it is preferably substituted by one, two or three, in particular one or two, identical or different residues  $R^4$ .  $R^4$  preferably is hydroxy,  $(C_1-C_4)$ -alkyloxy, di- $((C_1-C_4)$ -alkyl)amino or heteroaryl.

10  $R^5$  and  $R^6$  preferably are independently of one another hydrogen,  $(C_1-C_9)$ -alkyl,  $(C_1-C_4)$ -alkyl-O- $(C_1-C_3)$ -alkyl- or 5-membered or 6-membered aryl or together with the nitrogen atom carrying  $R^5$  and  $R^6$  form a 5-membered to 7-membered heterocycle which in addition to the nitrogen atom carrying the groups  $R^5$  and  $R^6$  can contain one further ring heteroatom from the series N, O and S and which can be substituted by one or more, for example one, two, three or four, identical or different residues from the series  $(C_1-$   
 15  $C_3)$ -alkyl, hydroxy- $(C_1-C_3)$ -alkyl-, 5-membered or 6-membered aryl, carbamoyl, hydroxy and oxo. It is particularly preferred if  $R^5$  and  $R^6$  together with the nitrogen atom carrying these residues form a 5-membered, 6-membered or 7-membered heterocycle which in addition to the nitrogen atom carrying the groups  $R^5$  and  $R^6$  can contain one further ring heteroatom from the series N, O and S and which can be substituted by one or more,  
 20 for example one, two, three or four, identical or different residues from the series  $(C_1-C_3)$ -alkyl, hydroxy- $(C_1-C_3)$ -alkyl-, 5-membered or 6-membered aryl, carbamoyl, hydroxy

and oxo, in particular (C<sub>1</sub>-C<sub>3</sub>)-alkyl such as, for example, methyl. Preferably a heterocycle which is formed from the groups R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom carrying these residues is saturated.

- 5 An especially preferable heterocycle which is formed by R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom carrying these residues, is derived from morpholine, thiomorpholine, 1,1-dioxo-thiomorpholine, 1-oxo-thiomorpholine, from dialkylmorpholines such as dimethylmorpholines, from 2,6-dimethylmorpholine, cis-2,6-dimethylmorpholine, 3,5-dimethylmorpholine, cis-3,5-dimethylmorpholine, 1-(pyrimidin-2-yl)-piperazine,  
10 piperidine-4-carboxamide, 1-(2-hydroxyethyl)-piperazine, 1-methylpiperazine, 1-ethylpiperazine, from 1-arylpiperazines, from ethyl piperazine-1-carboxylate, piperidine, 2-methylpiperidine, 2,6-dimethylpiperidine, cis-2,6-dimethylpiperidine, 3,5-dimethylpiperidine, cis-3,5-dimethylpiperidine, 4-hydroxypiperidine, from 4-oxopiperidine or a ketal thereof like 1,4-dioxo-8-aza-spiro[4.5]decan, from tetrahydropyridine,  
15 tetrahydropyrimidine, 1-methylhomopiperazine, thiazolidine, pyrroline, pyrrolidine, 3-hydroxypyrrolidine, 1,2,3,4-tetrahydroisoquinoline or 2,3-dihydro-1H-isoindole, where these rings are attached via the ring nitrogen atom or, in the case of piperazine derivatives, via the unsubstituted ring nitrogen atom. A more especially preferable heterocycle which is formed by R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom carrying these  
20 residues, is derived from morpholine, thiomorpholine, 1,1-dioxo-thiomorpholine, 1-oxo-thiomorpholine, 2,6-dimethylmorpholine, cis-2,6-dimethylmorpholine, 3,5-

dimethylmorpholine, cis-3,5-dimethylmorpholine, 1-(pyrimidin-2-yl)-piperazine, piperidine-4-carboxamide, 1,2,3,4-tetrahydroisoquinoline or 2,3-dihydro-1H-isoindole, moreover preferably from morpholine, 2,6-dimethylmorpholine or cis-2,6-dimethylmorpholine, in particular from morpholine or cis-2,6-dimethylmorpholine, where  
5 these rings are attached via the ring nitrogen atom or, in the case of the piperazine derivative, via the unsubstituted ring nitrogen atom.

R<sup>7</sup> preferably is hydrogen, (C<sub>1</sub>-C<sub>3</sub>)-alkyl, ((C<sub>1</sub>-C<sub>4</sub>)-alkyl)<sub>2</sub>N-(C<sub>1</sub>-C<sub>3</sub>)-alkyl- or (C<sub>1</sub>-C<sub>4</sub>)-alkyl-O-(C<sub>1</sub>-C<sub>3</sub>)-alkyl-.

R<sup>8</sup> preferably is hydrogen, (C<sub>1</sub>-C<sub>3</sub>)-alkyl or acetyl.

Aryl preferably is phenyl or heteroaryl, in particular phenyl or 5-membered or 6-membered heteroaryl. Unless stated otherwise, preferred substituents on aryl residues  
15 are halogen, CF<sub>3</sub>, (C<sub>1</sub>-C<sub>3</sub>)-alkyl, cyano, nitro and (C<sub>1</sub>-C<sub>3</sub>)-alkyloxy, more preferred substituents are CF<sub>3</sub>, chlorine, methyl and methoxy.

Heteroaryl and heteroarylene preferably are a residue of a monocyclic 5-membered or 6-membered aromatic heterocycle, in particular a residue derived from the  
20 heteroaromatics thiophene, pyrazole, thiazole, oxazole, isoxazole, pyridine, pyrimidine, pyridazine and tetrazole.



Heterocyclyl preferably is a residue derived from a saturated heterocycle, more preferably a residue of a monocyclic 5-membered or 6-membered saturated heterocycle, in particular a residue which is derived from pyrrolidine, piperidine, from N-alkylpiperazines, from morpholine, from dialkylmorpholines, from thiomorpholine or tetrahydrofuran. In addition, the above explanations on preferred heterocycles which are formed by the residues  $R^5$  and  $R^6$  together with the nitrogen atom carrying these residues correspondingly apply to heterocyclyl residues which are attached via a ring nitrogen atom.

If a group  $S(O)_n$  is bonded to a nitrogen atom the number  $n$  therein preferably is 1 or 2, more preferably 2. The number  $n$  in the group  $R^1-S(O)_n$  preferably is 0 or 2, particularly preferably 2.

X preferably is O or a nitrogen atom which via a single bond is attached to a ring carbon atom in the group  $A^1$  which ring carbon atom is directly adjacent to the carbon atom in  $A^1$  carrying the group  $-NH-C(=X)-$  so that the group  $-NH-C(=X)-$  together with the carbon atoms in  $A^1$  carrying it forms an anellated imidazole ring. Particularly preferably X is O.

Preferred compounds of the formula I are those compounds in which one or more of the residues contained therein have preferred meanings, all combinations of preferred

substituent definitions being a subject of the present invention. Also with respect to all preferred compounds of the formula I the present invention includes all stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts.

Groups of preferred compounds are also formed, for example, by the compounds of the formulae Ia, Ib, Ic, Id, Ie, If, Ig and Ih in which one or more residues have preferred meanings, in all their stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts.

A group of particularly preferred compounds is formed, for example, by compounds of the formula I in which

A<sup>1</sup> is phenylene or heteroarylene where these residues can be unsubstituted or substituted by one or more identical or different substituents from the series halogen, (C<sub>1</sub>-C<sub>4</sub>)-alkyl, CF<sub>3</sub>, -O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl and -CN;

the ring A<sup>2</sup> which comprises the two carbon atoms which carry the groups R<sup>2</sup>-SO<sub>2</sub>-NH and C(=X)-NH- is an aromatic ring;

R<sup>1</sup> is (C<sub>1</sub>-C<sub>7</sub>)-alkyl which can be substituted by one or more identical or different residues R<sup>4</sup>, or is aryl, or if the number n in the group R<sup>1</sup>-S(O)<sub>n</sub>- is 2 also is NR<sup>5</sup>R<sup>6</sup>;

R<sup>2</sup> is aryl;

R<sup>3</sup> denotes one or more identical or different residues from the series hydrogen, halogen, CF<sub>3</sub>, OH, -O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl, -O-(C<sub>2</sub>-C<sub>4</sub>)-alkyl-O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl, -O-aryl, NO<sub>2</sub>, -CN, NR<sup>7</sup>R<sup>8</sup>, -CO-NR<sup>7</sup>R<sup>8</sup>, -CO-OH, -CO-O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl, heterocyclyl, -S(O)<sub>n</sub>-(C<sub>1</sub>-C<sub>4</sub>)-alkyl and (C<sub>1</sub>-C<sub>4</sub>)-alkyl which can be substituted by one or more identical or different residues R<sup>4</sup>;

5

R<sup>4</sup> is fluorine, OH, -O-(C<sub>1</sub>-C<sub>10</sub>)-alkyl, -O-(C<sub>2</sub>-C<sub>4</sub>)-alkyl-O-(C<sub>1</sub>-C<sub>7</sub>)-alkyl, -O-aryl, -CN, NR<sup>7</sup>R<sup>8</sup>, -CO-NH<sub>2</sub>, -CO-NH-(C<sub>1</sub>-C<sub>3</sub>)-alkyl, -CO-N((C<sub>1</sub>-C<sub>3</sub>)-alkyl)<sub>2</sub>, -CO-OH, -CO-O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl, heterocyclyl or oxo;

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145

"aryl" is phenyl or heteroaryl which can all be substituted by one or more identical or different substituents from the series halogen, (C<sub>1</sub>-C<sub>4</sub>)-alkyl, phenyl, CF<sub>3</sub>, NO<sub>2</sub>, -O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl, (C<sub>1</sub>-C<sub>2</sub>)-alkylenedioxy, NH<sub>2</sub>, -NH-CO-(C<sub>1</sub>-C<sub>4</sub>)-alkyl, -CN, -CO-NH<sub>2</sub>, -CO-OH and -CO-O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl;

5

heteroaryl and heteroarylene are a residue of a monocyclic 5-membered or 6-membered aromatic heterocycle which contains one or more identical or different ring heteroatoms from the series N, O and S;

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185

A group of more particularly preferred compounds is formed, for example, by compounds of the formula I in which:

A<sup>1</sup> is phenylene which is unsubstituted or substituted by one or more identical or different substituents from the series halogen, (C<sub>1</sub>-C<sub>4</sub>)-alkyl, CF<sub>3</sub>, -O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl and  
5 -CN;

the ring A<sup>2</sup> which comprises the two carbon atoms which carry the groups R<sup>2</sup>-SO<sub>2</sub>-NH and C(=X)-NH- is a benzene ring;

10 R<sup>1</sup> is NR<sup>5</sup>R<sup>6</sup>;

R<sup>2</sup> is aryl;

R<sup>3</sup> denotes one or more identical or different residues from the series hydrogen,

15 halogen, CF<sub>3</sub>, -O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl, -CN and (C<sub>1</sub>-C<sub>4</sub>)-alkyl;

R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom carrying R<sup>5</sup> and R<sup>6</sup> form a 5-membered or 6-membered saturated heterocycle which in addition to the nitrogen atom carrying the groups R<sup>5</sup> and R<sup>6</sup> can contain one further ring heteroatom from the series N, O and S

20 and which can be substituted by one or more identical or different residues from the series (C<sub>1</sub>-C<sub>3</sub>)-alkyl, hydroxy-(C<sub>1</sub>-C<sub>3</sub>)-alkyl-, aryl, carbamoyl, hydroxy and oxo;

aryl is phenyl or 5-membered or 6-membered heteroaryl containing one or more identical or different ring heteroatoms from the series N, O and S which residues can all be substituted by one or more identical or different substituents from the series halogen, (C<sub>1</sub>-C<sub>4</sub>)-alkyl, CF<sub>3</sub>, NO<sub>2</sub>, -O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl, -NH-CO-(C<sub>1</sub>-C<sub>4</sub>)-alkyl and -CN;

5

n is 2;

X is oxygen;

in all their stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts.

A group of especially preferred compounds is formed, for example, by compounds of the formula I in which:

15 A<sup>1</sup> is an unsubstituted divalent phenylene residue;

the ring A<sup>2</sup> which comprises the two carbon atoms which carry the groups R<sup>2</sup>-SO<sub>2</sub>-NH and C(=X)-NH- is a benzene ring;

20 R<sup>1</sup> is NR<sup>5</sup>R<sup>6</sup>;

R<sup>2</sup> is aryl;

R<sup>3</sup> denotes one or more identical or different residues from the series hydrogen, halogen, -O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl and (C<sub>1</sub>-C<sub>4</sub>)-alkyl;

5

R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom carrying R<sup>5</sup> and R<sup>6</sup> form a saturated 6-membered heterocycle which in addition to the nitrogen atom carrying the groups R<sup>5</sup> and R<sup>6</sup> can contain one further ring heteroatom from the series N, O and S and which can be substituted by one or more identical or different residues from the series (C<sub>1</sub>-C<sub>3</sub>)-alkyl, aryl, oxo and carbamoyl;

aryl is phenyl or 5-membered or 6-membered heteroaryl containing one or more identical or different ring heteroatoms from the series N, O and S which residues can all be substituted by one or more identical or different substituents from the series halogen, (C<sub>1</sub>-C<sub>4</sub>)-alkyl, CF<sub>3</sub> and -O-(C<sub>1</sub>-C<sub>4</sub>)-alkyl;

n is 2;

X is oxygen;

20

in all their stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts.

A group of even more preferred compounds is formed, for example, by compounds of the formula I in which:

A<sup>1</sup> is an unsubstituted divalent 1,4-phenylene residue;

the ring A<sup>2</sup> which comprises the two carbon atoms which carry the groups R<sup>2</sup>-SO<sub>2</sub>-NH and C(=X)-NH-, together with the residues R<sup>3</sup>, is a benzene ring which carries one or two substituents from the series chlorine and methoxy;

R<sup>1</sup> is NR<sup>5</sup>R<sup>6</sup>;

R<sup>2</sup> is phenyl or thienyl which residues are all substituted by one or two chlorine atoms;

R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom carrying R<sup>5</sup> and R<sup>6</sup> form a saturated 6-membered heterocycle which in addition to the nitrogen atom carrying the groups R<sup>5</sup> and R<sup>6</sup> can contain one further ring heteroatom from the series O and S and which is unsubstituted or substituted by one or two methyl residues;

n is 2;



X is oxygen;

in all their stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts.

5

The presents invention also relates to processes for the preparation of the compounds of the formula I which are described in the following and by which the compounds of the invention are obtainable. According to Scheme 1 compounds of the invention can be obtained, for example, by first reacting an aminocarboxylic acid of the formula II with a sulfonyl chloride of the formula  $R^2-SO_2-Cl$  or a sulfonic acid anhydride in the presence of a base in a solvent like water, pyridine or an ether. Suitable bases are inorganic bases like sodium carbonate or organic bases like, for example, pyridine or triethylamine. The sulfonylaminocarboxylic acid of the formula III that is obtained can then be activated, for example by reaction with a chlorinating agent like phosphorus pentachloride, phosphorus oxychloride or thionyl chloride in an inert solvent, to give an acid chloride of the formula IV which is then reacted with an arylamine.

The activation of the carboxylic acid group in the compounds of the formula III can, however, also be carried out in a different manner, for example by one of the numerous methods for the formation of amide bonds in peptide chemistry which are well-known to


the skilled person, for example by conversion into a mixed anhydride or an activated ester or by using a carbodiimide like dicyclohexylcarbodiimide.

The reaction of the activated sulfonylaminocarboxylic acid with an arylamine is favorably  
 5 carried out in an inert solvent such as, for example, pyridine, tetrahydrofuran or toluene  
 in the absence or in the presence of an inert auxiliary base like, for example, a tertiary  
 amine or pyridine. If the arylamine that is employed in the reaction with the activated  
 carboxylic acid already contains the desired substituent  $R^1-S(O)_n$  then the reaction  
 directly provides the final compound of the formula I. Compounds of the formula I in  
 10 which the number  $n$  in the group  $R^1-S(O)_n$  is 1 or 2 can also be obtained by reacting an  
 activated carboxylic acid with a mercapto substituted arylamine of the formula  $R^1-S-A^1-$   
 $NH_2$  and then oxidizing the mercapto group in the compound of the formula V under  
 standard conditions, for example with a peroxide like hydrogen peroxide or a peracid  
 like 3-chloroperbenzoic acid or monoperoxyphthalic acid in a solvent like, for example,  
 15 methylene chloride or acetone. The activated carboxylic acids can also first be reacted  
 with arylamines of the formula  $A^1-NH_2$ . The resulting reaction product of the formula VI  
 can then be chlorosulfonated under standard conditions and the chlorosulfonyl group  
 can then be converted under standard conditions into the group  $R^1-SO_2$ , for example by  
 reaction with suitable amines in substance or in a solvent like N-methylpyrrolidone,  
 20 dimethylformamide, toluene or an ether, optionally in the presence of an auxiliary base.  
 In a similar manner the activated carboxylic acids can be reacted with

fluorosulfonylarylamines of the formula  $F-SO_2-A^1-NH_2$  and the fluorosulfonyl intermediates of the formula VII that are obtained can be converted under standard conditions into the compounds of the formula I according to the invention.


2000-07-27 10:00:00

(Scheme 1)




The compounds of the formula I according to the invention can furthermore be obtained by reacting the activated sulfonylaminocarboxylic acids, for example the acid chlorides of the formula IV shown in Scheme 1, with a mercaptoarylamine of the formula  $H_2N-A^1-SH$  which is unsubstituted on the sulfur atom. In a nucleophilic substitution reaction the product of the formula VIII that is obtained can subsequently be alkylated or arylated on the sulfur atom with an alkyl halogenide or an aryl halogenide or another reactive compound under standard conditions and, if desired, be oxidized on the sulfur to give the sulfoxide or the sulfone as explained above with respect to the compounds of the formula V (see Scheme 2).

(Scheme 2)



Compounds of the formula I can also be obtained, for example, by first activating a suitably substituted nitrocarboxylic acid of the formula IX, for example by converting it into the respective acid chloride of the formula X or by another procedure, and then reacting it with a substituted arylamine of the formula  $R^1-S(O)_n-A^1-NH_2$  analogously to the procedures described above (see Scheme 3). Here, too, as arylamines fluorosulfonylarylamines of the formula  $F-SO_2-A^1-NH_2$  can be employed, and in the N-(fluorosulfonylaryl)-carboxamides of the formula XI that are obtained the fluorosulfonyl group can be converted under standard conditions into a group  $R^1-SO_2$  according to the invention, for example with an amine of the formula  $HNR^5R^6$ .

(Scheme 3)



Before the nitro group is reduced to an amino group, in the nitro intermediates of the formula XII, use can be made of the activating effect of the nitro group on the ring A<sup>2</sup> and a suitable residue R<sup>3</sup>, for example a halogen atom, can be replaced with a different residue R<sup>3</sup> by reaction with a nucleophile, for example an amine. The reduction of the

5 nitro group to give an amino group can, for example, be carried out by catalytic hydrogenation in the presence of a noble metal catalyst or, preferably, in the presence of Raney nickel in a solvent like ethanol, glacial acetic acid or an ethanolic solution of hydrogen chloride, or it can be carried out by reduction with a base metal like zinc, tin or iron in the presence of an acid. The reduction can also be carried out, for example, with

10 tin(II) chloride or by reaction with sodium dithionite, favorably in a mixture of methanol, tetrahydrofuran and water as solvent. The sulfonylation of the amino group in the reduction product of the formula XIII with an activated sulfonic acid derivative can be carried out analogously to the reactions described above, for example with a sulfonic acid chloride in pyridine, and finally gives the compound of the formula I.

15


The compounds of the formula I in which X is a nitrogen atom which via a single bond is attached to a ring carbon atom in the group A<sup>1</sup> which ring carbon atom is directly adjacent to the carbon atom in A<sup>1</sup> carrying the group -NH-C(=X)-, i. e. the benzimidazole derivatives of the formula II, can for example be obtained by reacting an

20 activated sulfonylaminocarboxylic acid derivative obtained as described above according to Scheme 1, for example a carboxylic acid chloride of the formula IV, (or



also, in analogy to Scheme 3, a nitrocarboxylic acid derivative) with a 1,2-diaminoarene in the presence of a dehydrating agent such as, for example, thionyl chloride or phosphorus pentachloride (see Scheme 4). The reaction is usually carried out in an inert solvent, for example in a hydrocarbon like toluene or xylene. The 1,2-diaminoarene can  
5 already contain the final group  $R^1-S(O)_n$  or a precursor group thereof, for example the group  $R^1-S$ . Subsequent steps, for example reactions on the sulfur atom, can then be carried out as explained above. Just so, unsubstituted 1,2-diaminoarenes can be employed and the resulting products of the formula XIV can be chlorosulfonated, for example with chlorosulfuric acid, and the sulfonyl chlorides that are obtained can be  
10 converted into the final compounds containing the group  $R^1-SO_2$ , for example by reaction with a suitable amine.

(Scheme 4)



All reactions for the synthesis of the compounds of the formula I are per se well-known to the skilled person and can be carried out under standard conditions according to or analogously to procedures described in the literature, for example in Houben-Weyl, Methoden der Organischen Chemie (Methods of Organic Chemistry), Thieme-Verlag, Stuttgart, or Organic Reactions, John Wiley & Sons, New York. Depending on the circumstances of the individual case, in order to avoid side reactions during the synthesis of a compound of the formula I it can be necessary or advantageous to temporarily block functional groups by introducing protective groups and to deprotect them in a later stage of the synthesis, or introduce functional groups in the form of precursor groups which in a later reaction step are converted into the desired functional groups. Such synthetic strategies and protective groups and precursor groups which are suitable in an individual case are known to the skilled person. If desired the compounds of the formula I can be purified by customary purification procedures, for example by recrystallization or chromatography. The starting compounds for the preparation of the compounds of the formula I are commercially available or can be prepared according to or analogously to literature procedures.

The compounds of the formula I according to the invention effect an increase of the cGMP concentration via the activation of the soluble guanylate cyclase (sGC), and they are therefore useful agents for the therapy and prophylaxis of disorders which are associated with a low or decreased cGMP level or which are caused thereby, or for

whose therapy or prophylaxis an increase of the present cGMP level is desired. The activation of the sGC by the compounds of the formula I can be examined, for example, in the activity assay described below.

5 Disorders and pathological conditions which are associated with a low cGMP level or in which an increase of the cGMP level is desired and for whose therapy and prophylaxis it is possible to use compounds of the formula I are, for example, cardiovascular diseases, such as endothelial dysfunction, diastolic dysfunction, atherosclerosis, hypertension, stable and unstable angina pectoris, thromboses, restenoses, myocardial  
10 infarction, strokes, cardiac insufficiency or pulmonary hypertonia, or, for example, erectile dysfunction, asthma bronchiale, chronic kidney insufficiency and diabetes. Compounds of the formula I can additionally be used in the therapy of cirrhosis of the liver and also for improving a restricted memory performance or ability to learn.

15 The compounds of the formula I and their physiologically acceptable salts can be administered to animals, preferably to mammals, and in particular to humans, as pharmaceuticals by themselves, in mixtures with one another or in the form of pharmaceutical preparations. A subject of the present invention therefore also are the compounds of the formula I and their physiologically acceptable salts for use as  
20 pharmaceuticals, their use for activating soluble guanylate cyclase, for normalizing a disturbed cGMP balance and in particular their use in the therapy and prophylaxis of the

abovementioned syndromes as well as their use for preparing medicaments for these purposes.

Furthermore, a subject of the present invention are pharmaceutical preparations (or  
5 pharmaceutical compositions) which comprise as active component an effective dose of  
at least one compound of the formula I and/or a physiologically acceptable salt thereof  
and a customary pharmaceutically acceptable carrier, i. e. one or more  
pharmaceutically acceptable carrier substances and/or additives. A subject of the  
present invention is also those compounds of the formula I which were already known  
40 per se and which are excluded by disclaimer from the above-defined compounds of the  
formula I which are per se a subject of the present invention, and their physiologically  
acceptable salt as activators of soluble guanylate cyclase.

All statements above and below relating to the pharmacological effects and the uses of  
15 the compounds of the formula I thus also apply to the compound of the formula I  
wherein simultaneously the ring  $A^2$  which comprises the carbon atoms which carry the  
groups  $C(=X)-NH-$  and  $NH-SO_2R^2$  is a benzene ring which is substituted in positions 3  
and 5 by chlorine,  $R^2$  is methyl, X is oxygen and  $R^1-S(O)_n-A^1-$  is a 5-chloro-2-(4-  
chlorophenylmercapto)-phenyl residue, and its physiologically acceptable salts.

Thus, a subject of the invention are, for example, said compound and its physiologically acceptable salts for use as a pharmaceutical, pharmaceutical preparations which comprise as active component an effective dose of said compound and/or a physiologically acceptable salt thereof and a customary pharmaceutically acceptable carrier, and the uses of said compound and/or a physiologically acceptable salt thereof in the therapy or prophylaxis of the abovementioned syndromes as well as their use for preparing medicaments for these purposes.

The pharmaceuticals according to the invention can be administered orally, for example in the form of pills, tablets, lacquered tablets, sugar-coated tablets, granules, hard and soft gelatin capsules, aqueous, alcoholic or oily solutions, syrups, emulsions or suspensions, or rectally, for example in the form of suppositories. Administration can also be carried out parenterally, for example subcutaneously, intramuscularly or intravenously in the form of solutions for injection or infusion. Other suitable administration forms are, for example, percutaneous or topical administration, for example in the form of ointments, tinctures, sprays or transdermal therapeutic systems, or the inhalative administration in the form of nasal sprays or aerosol mixtures, or, for example, microcapsules, implants or rods. The preferred administration form depends, for example, on the disease to be treated and on its severity.

The amount of active compound of the formula I and/or its physiologically acceptable salts in the pharmaceutical preparations normally is from 0.2 to 500 mg, preferably from 1 to 200 mg, per dose, but depending on the type of the pharmaceutical preparation it can also be higher. The pharmaceutical preparations usually comprise 0.5 to 90 percent  
5 by weight of the compounds of the formula I and/or their physiologically acceptable salts. The preparation of the pharmaceutical preparations can be carried out in a manner known per se. For this purpose, one or more compounds of the formula I and/or their physiologically acceptable salts, together with one or more solid or liquid pharmaceutical carrier substances and/or additives (or auxiliary substances) and, if  
10 desired, in combination with other pharmaceutically active compounds having therapeutic or prophylactic action, are brought into a suitable administration form or dosage form which can then be used as a pharmaceutical in human or veterinary medicine.

15 For the production of pills, tablets, sugar-coated tablets and hard gelatin capsules it is possible to use, for example, lactose, starch, for example maize starch, or starch derivatives, talc, stearic acid or its salts, etc. Carriers for soft gelatin capsules and suppositories are, for example, fats, waxes, semisolid and liquid polyols, natural or hardened oils, etc. Suitable carriers for the preparation of solutions, for example of  
20 solutions for injection, or of emulsions or syrups are, for example, water, physiologically sodium chloride solution, alcohols such as ethanol, glycerol, polyols, sucrose, invert

sugar, glucose, mannitol, vegetable oils, etc. It is also possible to lyophilize the compounds of the formula I and their physiologically acceptable salts and to use the resulting lyophilisates, for example, for preparing preparations for injection or infusion. Suitable carriers for microcapsules, implants or rods are, for example, copolymers of

5 glycolic acid and lactic acid.

Besides the active compounds and carriers, the pharmaceutical preparations can also contain customary additives, for example fillers, disintegrants, binders, lubricants, wetting agents, stabilizers, emulsifiers, dispersants, preservatives, sweeteners,

10 colorants, flavorings, aromatizers, thickeners, diluents, buffer substances, solvents, solubilizers, agents for achieving a depot effect, salts for altering the osmotic pressure, coating agents or antioxidants.

The dosage of the active compound of the formula I to be administered and/or of a

15 physiologically acceptable salt thereof depends on the individual case and is, as is customary, to be adapted to the individual circumstances to achieve an optimum effect. Thus, it depends on the nature and the severity of the disorder to be treated, and also on the sex, age, weight and individual responsiveness of the human or animal to be treated, on the efficacy and duration of action of the compounds used, on whether the

20 therapy is acute or chronic or prophylactic, or on whether other active compounds are administered in addition to compounds of the formula I. In general, a daily dose of



approximately 0.01 to 100 mg/kg, preferably 0.1 to 10 mg/kg, in particular 0.3 to 5 mg/kg (in each case mg per kg of bodyweight) is appropriate for administration to an adult weighing approximately 75 kg in order to obtain the desired results. The daily dose can be administered in a single dose or, in particular when larger amounts are administered, be divided into several, for example two, three or four individual doses. In some cases, depending on the individual response, it may be necessary to deviate upwards or downwards from the given daily dose.

The compounds of the formula I activate the soluble guanylate cyclase, mainly by binding in the heme binding pocket of the enzyme. On account of this property, apart from use as pharmaceutically active compounds in human medicine and veterinary medicine, they can also be employed as a scientific tool or as aids for biochemical investigations in which such an effect on guanylate cyclase is intended, and also for diagnostic purposes, for example in the in vitro diagnosis of cell samples or tissue samples. The compounds of the formula I and salts thereof can furthermore be employed, as already mentioned above, as intermediates for the preparation of other pharmaceutically active compounds.

The following examples of compounds of the formula I and of intermediates for their preparation illustrate the invention without limiting it.

## Examples

### 1.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoic acid

5 33.71 g (0.32 mol) of sodium carbonate were dissolved in 250 ml of water and warmed to 60 °C. 25.00 g (0.13 mol) of 2-amino-4,5-dimethoxy-benzoic acid were introduced into the solution, and to this solution 29.55 g (0.14 mol) of 4-chloro-benzenesulfonyl chloride were added portionwise over 15 min . After cooling the mixture was filtered with suction, the residue was taken up in 1 % sodium hydrogencarbonate solution, the  
10 solution filtered, and the product precipitated by addition of 1 N hydrochloric acid. 25.90 g (55 %) of 2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoic acid of melting point (m.p.) 212-214 °C were obtained.

Analogously were obtained:

15

2.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoic acid; M.p.: 210 °C

3.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-benzoic acid

20 5.) 2-(4-Chloro-phenylsulfonylamino)-cyclopentanecarboxylic acid; M.p.: 147 °C

6.) 2-(4-Chloro-phenylsulfonylamino)-5-methyl-benzoic acid; M.p.: 201 °C

7.) 3-(4-Chloro-phenylsulfonylamino)-thiophene-2-carboxylic acid; M.p.: 180 °C

5 8.) 3-(4-Chloro-phenylsulfonylamino)-pyrazole-4-carboxylic acid; Oil

9.) 2-(4-Chloro-phenylsulfonylamino)-pyridine-3-carboxylic acid; M.p.: Decomposition  
(dec.) > 360 °C

10 10.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl chloride

25.90 g (0.07 mol) of 2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoic acid were  
mixed with 75 ml of toluene, 17.30 g (0.08 mol) of phosphorus pentachloride were  
added and the mixture was stirred at 40-45 °C for 2.5 h. Then the mixture was  
15 concentrated in vacuo to half of its volume and the product that precipitated was filtered  
off with suction and washed with a small amount of toluene. 25.30 g (93 %) of 2-(4-  
chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl chloride having a melting point of  
175-177 °C were obtained.

20 Analogously were obtained:

11.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoyl chloride; M.p.: 127 °C

12.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-benzoyl chloride; M.p.: 117 °C

5 13.) 2-(4-Chloro-phenylsulfonylamino)-cyclopentanecarboxylic acid chloride; M.p.: 107 °C

14.) 2-(4-Chloro-phenylsulfonylamino)-5-methyl-benzoyl chloride; M.p.: 114 °C

10 15.) 3-(4-Chloro-phenylsulfonylamino)-thiophene-2-carboxylic acid chloride; M.p.: 122 °C

16.) 3-(4-Chloro-phenylsulfonylamino)-pyrazole-4-carboxylic acid chloride; M.p.: 260 °C  
(Dec.)

15

17.) 2-(4-Chloro-phenylsulfonylamino)-pyridine-3-carboxylic acid chloride

18.) 4-((2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl)-amino)-  
benzenesulfonyl fluoride

20

10.00 g (25.6 mmol) of 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl chloride were mixed with 300 ml of toluene, 4.49 g (25.6 mmol) of 4-aminobenzenesulfonyl fluoride were added and the mixture was heated under reflux for 4 h. After cooling the precipitated solid was filtered off with suction and washed with 5 toluene. 11.71 g (87 %) of the title compound having a melting point of 216-219 °C were obtained.

Analogously were obtained:

19.) 4-((5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoyl)-amino)-benzenesulfonyl fluoride; M.p.: 242 °C

20.) N-(4-Aminosulfonyl-phenyl)-5-chloro-2-(4-chloro-phenylsulfonylamino)-benzamide; M.p.: 260 °C

21.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-((4-(4-nitro-phenyl)-mercapto)-phenyl)-benzamide; M.p.: 255 °C

22.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(phenylmercapto)-phenyl)-benzamide; M.p.: 169 °C

23.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-methylmercapto-phenyl)-benzamide; M.p.: 220 °C

24.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-methyl-benzothiazol-5-yl)-

5 benzamide; M.p.: 251 °C

25.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(3-diethylamino-2-hydroxy-propylmercapto)-phenyl)-benzamide; M.p.: 102 °C

10 26.) 4-((5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-benzoyl)-amino)-benzenesulfonyl fluoride; M.p.: 232 °C

27.) 4-(2-(4-Chloro-phenylsulfonylamino)-cyclopentanecarbonylamino)-benzenesulfonyl fluoride; M.p.: 211 °C

15

28.) 4-((2-(4-Chloro-phenylsulfonylamino)-5-methyl-benzoyl)-amino)-benzenesulfonyl fluoride; M.p.: 224 °C

29.) 4-((3-(4-Chloro-phenylsulfonylamino)-thiophene-2-carbonyl)-amino)-

20 benzenesulfonyl fluoride; M.p.: 255 °C

30.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-mercapto-phenyl)-benzamide;

M.p.: 202 °C

31.) 4-((3-(4-Chloro-phenylsulfonylamino)-pyrazol-4-carbonyl)-amino)-benzenesulfonyl

5 fluoride; M.p.: 251 °C

32.) 3-((5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoyl)-amino)-benzenesulfonyl

fluoride; M.p.: 224 °C

33.) 4-(2-(4-Chloro-phenylsulfonylamino)-pyridine-3-carbonyl)-amino)-benzenesulfonyl

fluoride; M.p.: 263-265 °C

34.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-methyl-5-(thiomorpholine-4-

sulfonyl)-thiazol-2-yl)-benzamide; M.p.: 265-267 °C

35.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-methylmercapto-phenyl)-

benzamide

36.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-methylmercapto-phenyl)-

benzamide

37.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(5-methyl-isoxazol-3-yl-sulfamoyl)-phenyl)-benzamide

38.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-nitro-phenylsulfonyl)-phenyl)-

5 benzamide

39.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(5-ethylsulfonyl-2-hydroxy-phenyl)-benzamide

40.) N-(3-Butylsulfamoyl-phenyl)-5-chloro-2-(4-chloro-phenylsulfonylamino)-benzamide

41.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-nitro-5-propylmercapto-phenyl)-benzamide

42.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-thiocyanato-phenyl)-benzamide

43.) N-(4-Acetylsulfamoyl-phenyl)-5-chloro-2-(4-chloro-phenylsulfonylamino)-benzamide

44.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-phenylmercapto-phenyl)-benzamide



45.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-chloro-5-(2-cyano-ethylsulfamoyl)-phenyl)-benzamide

46.) N-(5-Butylsulfamoyl-2-methoxy-phenyl)-5-chloro-2-(4-chloro-phenylsulfonylamino)-  
5 benzamide

47.) N-(4-Benzoylsulfamoyl-phenyl)-5-chloro-2-(4-chloro-phenylsulfonylamino)-  
benzamide

48.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-chloro-4-methylsulfonyl-phenyl)-  
benzamide

49.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(hexadecylsulfonyl)-phenyl)-  
benzamide

50.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-  
(butylaminocarbonylamino-sulfonyl)-phenyl)-benzamide

51.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-sulfamoyl-phenyl)-benzamide

52.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-methylmercapto-5-trifluoromethyl-phenyl)-benzamide

53.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-methylsulfonyl-phenyl)-benzamide

5

54.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-(2-hydroxy-ethylsulfonyl)-phenyl)-benzamide

55.) (4-(5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoylamino)-phenylmercapto)-acetic acid

10

56.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(3,4-dimethyl-isoxazol-5-ylsulfamoyl)-phenyl)-benzamide

15 57.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(thiazol-2-ylsulfamoyl)-phenyl)-benzamide

58.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-N-(4-ethylmercapto-phenyl)-benzamide; M.p.: 171 °C

20

59.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide

500 mg (0.95 mmol) of 4-((2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl)-amino)-benzenesulfonyl fluoride were dissolved in 1 ml of thiomorpholine and heated to 90 °C for 30 min. For working up the mixture was poured onto 50 ml of ice/1 N hydrochloric acid, the precipitate was filtered off with suction, dried in a vacuum-drying chamber over phosphorus pentoxide and recrystallized from hexane/ethyl acetate. 378 mg (65 %) of the title compound having a melting point of 241 °C were obtained.

Analogously were obtained:

60.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-pyridine-3-carboxamide; M.p.: 256-258 °C

61.) N-(4-(4-Carbamoyl-piperidine-1-sulfonyl)-phenyl)-2-(4-chloro-phenylsulfonylamino)-pyridine-3-carboxamide; M.p.: 273-276 °C

62.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(piperidine-1-sulfonyl)-phenyl)-pyridine-3-carboxamide; M.p.: 180-183 °C

63.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 246 °C

64.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-methyl-piperazine-1-sulfonyl)-phenyl)-benzamide; M.p.: 219 °C

65.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 259 °C

66.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 251 °C

67.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-hydroxy-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 255 °C

68.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(1,4-dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-phenyl)-benzamide; M.p.: 256 °C

69.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 253 °C

70.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 222 °C

71.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-N-(4-(4-methyl-piperazine-1-sulfonyl)-phenyl)-benzamide; M.p.: 246 °C

72.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 172 °C

73.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-(2-hydroxy-ethyl)-piperazine-1-sulfonyl)-phenyl)-benzamide; M.p.: 277 °C

74.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-cyclopentanecarboxamide; M.p.: 180 °C

75.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-diethylsulfamoyl-phenyl)-benzamide; M.p.: 226 °C

76.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 240 °C

77.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(2-methoxy-ethylsulfamoyl)-phenyl)-benzamide; M.p.: 209 °C

78.) 2-(4-Chloro-phenylsulfonylamino)-5-methyl-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 203 °C

79.) 3-(4-Chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-thiophene-2-carboxamide; M.p.: 220 °C

80.) 3-(4-Chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-1H-pyrazole-4-carboxamide; Oil

81.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 238 °C

82.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 202 °C

83.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-(4-methyl-piperazine-1-sulfonyl)-phenyl)-benzamide hydrochloride; M.p.: 245 °C

84.) 3-(4-Chloro-phenylsulfonylamino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-  
thiophene-2-carboxamide; M.p.: 229 °C

85.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-

5 benzamide; M.p.: 228 °C

86.) 2-(4-Chloro-phenylsulfonylamino)-5-methyl-N-(4-(thiomorpholine-4-sulfonyl)-  
phenyl)-benzamide; M.p.: 234 °C

87.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(4-methyl-piperazine-1-  
sulfonyl)-phenyl)-benzamide; M.p.: 172 °C

88.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-  
phenyl)-4,5-dimethoxy-benzamide; M.p.: 208 °C

89.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(4-hydroxy-piperidine-1-sulfonyl)-phenyl)-  
4,5-dimethoxy-benzamide; M.p.: 244 °C

90.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(piperidine-3-sulfonyl)-

phenyl)-benzamide; M.p.: 258 °C

91.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(thiazolidine-3-sulfonyl)-phenyl)-benzamide; M.p.: 261 °C

92.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(2,5-dihydro-1H-pyrrole-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 262 °C

93.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(1,2,3,6-tetrahydro-pyridine-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 252 °C

94.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(2-methyl-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 227 °C

95.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(piperazine-1-sulfonyl)-phenyl)-benzamide; M.p.: 243 °C

96.) 4-(4-(2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoylamino)-phenylsulfonyl)-piperazine-1-carboxylic acid ethyl ester; M.p.: 245 °C

97.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(4-methyl-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 267 °C



98.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(4-methyl-perhydro-[1,4]diazepine-1-sulfonyl)-phenyl)-benzamide; M.p.: 274 °C

99.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(4-ethyl-piperazine-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 191 °C

100.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-((2-dimethylamino-ethyl)-ethyl-sulfamoyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: Dec. > 119 °C

101.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(1,4,5,6-tetrahydro-pyrimidine-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: Dec. > 237 °C

102.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(4-(pyrimidin-2-yl)-piperazine-1-sulfonyl)-phenyl)-benzamide; M.p.: Dec. > 194 °C

103.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(4-(4-chloro-phenyl)-piperazine-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: Dec. > 243 °C

104.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(indan-1-ylsulfamoyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 161 °C

105.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-((2-(1H-indol-3-yl)-ethyl)-methyl-sulfamoyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 182 °C

106.) 1-(4-((2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl)-amino)-phenylsulfonyl)-piperidine-4-carboxamide; M.p.: 252 °C

107.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-cyclopropylsulfamoyl-phenyl)-4,5-dimethoxy-benzamide; M.p.: 222 °C

108.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(3-hydroxy-pyrrolidine-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 272 °C

109.) N-(4-(Allyl-cyclohexyl-sulfamoyl)-phenyl)-2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-benzamide; M.p.: 182 °C

110.) 1-(4-((2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl)-amino)-phenylsulfonyl)-pyrrolidine-2-carboxylic acid; M.p.: 240 °C (sintering)

111.) 5-Chloro-2-nitro-benzoyl chloride

100.00 g (0.50 mol) of 5-chloro-2-nitrobenzoic acid were mixed with 72.20 g (0.61 mol) of thionyl chloride and the mixture was heated under reflux for 2 h. The excess thionyl chloride was removed in vacuo. 106.50 g (ca. 98 %) of crude 5-chloro-2-nitro-benzoyl chloride were obtained as an oil.

5

Analogously was obtained:

112.) 5-Methyl-2-nitro-benzoyl chloride; Oil

113.) 4-(5-Chloro-2-nitro-benzoylamino)-benzenesulfonyl fluoride

86.00 g (0.39 mol) of 5-chloro-2-nitro-benzoyl chloride was dissolved in 300 ml of toluene, a solution of 62.00 g (0.35 mol) of 4-aminobenzenesulfonyl fluoride was added dropwise, and the mixture was heated under reflux for 4 h. Subsequently it was cooled, concentrated in vacuo to half of its volume, cooled, and the precipitated solid was filtered off with suction. 121.60 g (86 %) of the title compound having a melting point of 182-184 °C were obtained.

Analogously were obtained:

114.) 4-(5-Methyl-2-nitro-benzoylamino)-benzenesulfonyl fluoride; M.p.: 179 °C

115.) 5-Chloro-N-(4-ethylmercapto-phenyl)-2-nitro-benzamide

116.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-nitro-benzamide

5 120.00 g (0.33 mol) of 4-(5-chloro-2-nitro-benzoylamino)-benzenesulfonyl fluoride, 29.10 g (0.33 mol) of morpholine and 33.85 g (0.33 mol) of triethylamine were stirred in 1200 ml of toluene at 60 °C for 2 days. The precipitated solid was filtered off with suction and recrystallized from acetone/n-hexane. 102.10 g (71 %) of the title compound having a melting point of 243-245 °C were obtained.

Analogously were obtained:

117.) 5-Chloro-2-nitro-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 120 °C

118.) 5-Methyl-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-nitro-benzamide; M.p.: 249 °C

119.) N-(4-(Morpholine-4-sulfonyl)-phenyl)-5-(morpholin-4-yl)-2-nitro-benzamide

20 20.00 g (0.56 mol) of 4-(5-chloro-2-nitro-benzoylamino)-benzenesulfonyl fluoride in 48.5 g (0.557 mol) of morpholine were heated under reflux for 1 h. Subsequently the mixture

was cooled, poured onto ice/hydrochloric acid and filtered with suction. 26.0 g (98 %) of the title compound having a melting point of 252 °C were obtained.

120.) 2-Amino-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide

5

11.10 g (26.1 mmol) of 5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-nitro-benzamide were dissolved in 440 ml of tetrahydrofuran/methanol (1:1) and a solution of 27.23 g (156.4 mmol) of sodium dithionite in 330 ml of water was added dropwise. After stirring for 1 h at room temperature the organic solvents were removed in rotary evaporator, and the precipitated product was filtered off with suction and purified by chromatography over silica with methylene chloride/methanol (9:1). 5.68 g (55 %) of the title compound having a melting point of 229-231 °C were obtained.

Analogously were obtained:

15

121.) 2-Amino-5-chloro-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 177 °C

122.) 2-Amino-N-(4-(morpholine-4-sulfonyl)-phenyl)-(5-morpholin-4-yl)-benzamide;

20 M.p.: 228 °C

123.) 2-Amino-5-chloro-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: 159-161 °C

124.) 5-Chloro-2-(5-chloro-1,3-dimethyl-1H-pyrazole-4-sulfonyl-amino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide

5

250 mg (0.60 mmol) of 2-amino-5-chloro-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide were dissolved in 10 ml of dry pyridine, and a solution of 195 mg (0.85 mmol) 5-chloro-1,3-dimethyl-1H-pyrazole-4-sulfonyl chloride in 5 ml of pyridine was added dropwise at 0 °C. After 2 h the mixture was poured onto ice, the precipitated solid was filtered off with suction and purified by chromatography over silica with methylene chloride/methanol (98:2). 250 mg (69 %) of the title compound having a melting point of 215-216 °C were obtained.

Analogously were obtained:

15

125.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(4-methyl-phenylsulfonylamino)-benzamide; M.p.: 214 °C

20

126.) 5-Chloro-2-(3,4-dimethoxy-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 245 °C

127.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(4-trifluoromethoxy-phenylsulfonylamino)-benzamide; M.p.: 195 °C

128.) 2-((4-Acetylamino-phenyl)-sulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 198 °C

129.) 5-Chloro-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 112 °C

130.) 5-Chloro-2-(5-chloro-1,3-dimethyl-pyrazole-4-sulfonyl-amino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 161 °C

131.) 5-Chloro-2-((1-methyl-imidazole-4-sulfonyl)-amino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 141 °C

132.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(pyridine-3-sulfonylamino)-benzamide; M.p.: 222 °C

133.) 2-(4-Benzoyloxy-phenylsulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 245 °C

134.) 5-Chloro-2-(ethylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide;  
M.p.: 274-276 °C

135.) 2-((2-Acetamideo-4-methyl-thiazole-5-sulfonyl)-amino)-5-chloro-N-(4-(morpholine-  
5 4-sulfonyl)-phenyl)-benzamide; M.p.: 257 °C

136.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(thiophene-2-sulfonylamino)-  
benzamide; M.p.: 216 °C

137.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(4-trifluoromethyl-  
phenylsulfonylamino)-benzamide; M.p.: 264 °C

138.) 2-(4-Bromo-phenylsulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-  
benzamide; M.p.: 232 °C

139.) 2-(3,5-Bis-trifluoromethyl-phenylsulfonylamino)-5-chloro-N-(4-(morpholine-4-  
sulfonyl)-phenyl)-benzamide; M.p.: 209 °C

140.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(4-nitro-phenylsulfonylamino)-  
20 benzamide; M.p.: 239 °C



141.) 5-Chloro-2-(4-cyano-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 238 °C

142.) 5-Chloro-2-(4-methylsulfonyl-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 181 °C

143.) 5-Chloro-2-(4-isopropyl-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 105 °C

144.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-((2-phenyl-ethenyl)-sulfonylamino)-benzamide; M.p.: 278 °C

145.) 5-Chloro-2-(4,5-dibromo-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 232 °C

146.) 5-Chloro-2-(4-fluoro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 245 °C

147.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(5-phenylsulfonyl-thiophene-2-sulfonylamino)-benzamide; M.p.: 103 °C

148.) 5-Chloro-2-(3-chloro-4-methoxy-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 274 °C

149.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(quinoline-8-sulfonylamino)-

5 benzamide; M.p.: 262 °C

150.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(2,4,6-trimethyl-phenylsulfonylamino)-benzamide; M.p.: 240 °C

10 151.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(3-nitro-phenylsulfonylamino)-benzamide; M.p.: 220 °C

152.) 5-Chloro-2-(4-methoxy-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 269 °C

15

153.) 5-Chloro-2-methylsulfonylamino-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 248 °C

154.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-phenylmethylsulfonylamino-

20 benzamide; M.p.: 106 °C

155.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(2,2,2-trifluoro-ethylsulfonylamino)-benzamide; M.p.: 208 °C

156.) 2-(Butyl-sulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide;

5 M.p.: 102 °C

157.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(3-trifluoromethyl-phenylsulfonylamino)-benzamide; M.p.: 212 °C

158.) 2-(4-Bromo-2,5-dichloro-thiophene-3-sulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 267 °C

159.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(2-trifluoromethyl-phenylsulfonylamino)-benzamide; M.p.: 234 °C

160.) 5-Chloro-2-(3-chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 206 °C

161.) 2-(4-Bromo-2-methoxy-phenylsulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 260 °C

162.) 5-Chloro-2-(2,6-dichloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 244 °C

163.) 5-Chloro-2-(2-cyano-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-

5 benzamide; M.p.: 200 °C

164.) 2-(4-Butoxy-phenylsulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 225 °C

165.) 5-Chloro-2-(7,7-dimethyl-2-oxo-bicyclo[2.2.1]heptane-1-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 120 °C

166.) 5-Chloro-2-(3-fluoro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 204 °C

167.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-5-(morpholin-4-yl)-benzamide; M.p.: 264 °C

168.) 5-Chloro-N-(4-ethylsulfonyl-phenyl)-2-(4-methyl-phenylsulfonylamino)-benzamide; M.p.: 188-192 °C

169.) 5-Chloro-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: 195-197 °C

170.) 5-Chloro-2-(4-chloro-3-nitro-phenylsulfonylamino)-N-(4-ethylsulfonyl-phenyl)-

5 benzamide; M.p.: 196-198 °C

171.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: 180-185 °C

172.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: Dec. > 249 °C

173.) 5-Chloro-2-ethylsulfonylamino-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: 103 °C

174.) 4-Chloro-N-(2-(1H-benzimidazol-2-yl)-4-chloro-phenyl)-benzenesulfonamide

1.00 g (2.7 mmol) of 5-chloro-2-(4-chloro-phenylsulfonylamino)-benzoyl chloride and 296 mg (2.7 mmol) of o-phenylenediamine in 150 ml of toluene were heated under reflux for 1 h. A small amount of solid was filtered off with suction and the filtrate was evaporated. The residue was taken up in 50 ml of toluene, 600 mg of thionyl chloride

were added and the mixture was again heated under reflux for 10 h. Subsequently it was cooled and the precipitated solid was filtered off with suction. 280 mg (25 %) of the title compound having a melting point of 225-228 °C were obtained.

5 175.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(1,1-dioxo-thiomorpholine-4-sulfonyl)-phenyl)-benzamide

and

176.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(1-oxo-thiomorpholine-4-sulfonyl)-phenyl)-benzamide

500 mg (0.82 mmol) of 2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide in 50 ml of acetone were cooled to 0 °C.

A solution of 371 mg (1.23 mmol) of 57 % m-chloroperbenzoic acid in 20 ml of acetone was added and the mixture was stirred at room temperature over night. For working up

15 it was poured onto water/ice and the precipitate was filtered off with suction. The two products obtained as a mixture were separated by chromatography over silica with methylene chloride/methanol (97:3).

Analogously were obtained:

177.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(1,1-dioxo-thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 182 °C

178.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(1-oxo-thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 233 °C

179.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: 240 °C

180.) 5-Chloro-N-(4-ethylsulfonyl-phenyl)-2-nitro-benzamide

181.) 4-Chloro-N-(4-chloro-2-(6-(morpholine-4-sulfonyl)-1H-benzimidazol-2-yl)-phenyl)-benzenesulfonamide

200 mg (0.5 mmol) of 4-chloro-N-(2-(1H-benzimidazol-2-yl)-4-chloro-phenyl)-benzenesulfonamide were added at 0 °C to 1 ml of chlorosulfuric acid and heated to 60 °C for 30 min. Subsequently the mixture was poured onto water/ice, and the solid was filtered off with suction, dried and added at 0 °C to 1 ml of morpholine. After stirring at room temperature for 1 h the mixture was poured onto ice/hydrochloric acid and extracted with ethyl acetate. The extracts were evaporated and the residue was purified

by chromatography over silica with hexane/ethyl acetate (1:1). 20 mg (7 %) of the title compound having a melting point of 225-228 °C were obtained.

<sup>1</sup>H-NMR (D<sub>6</sub>-DMSO): δ (ppm) = 2.9 (m, 4 H, morpholine-H), 3.6 (m, 4 H, morpholine-H), 7.5 (dd, 4 H, phenylene-H), 7.4-8.2 (m, 6 H, benzo-H, phenyl-H)

5

182.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(2-(pyrrolidin-1-yl)-ethylsulfonylamino)-benzamide

The compound was prepared by using 2-chloro-ethylsulfonyl chloride. The 1-(2-(4-chloro-2-(4-(morpholine-4-sulfonyl)-phenylcarbamoyl)-phenylsulfamoyl)-ethyl)-pyridinium chloride that was isolated as an intermediate was reacted with pyrrolidine to give the title compound.

<sup>1</sup>H-NMR (D<sub>6</sub>-DMSO): δ (ppm) = 1.8 (m, 4 H, pyrrolidine-H), 2.65 (m, 4 H, pyrrolidine-H), 3.0 (m, 4 H, morpholine-H), 3.1 (t, 2 H, ethylene-H), 3.35 (t, 2 H, ethylene-H), 3.75 (m, 4 H, morpholine-H), 7.50 (dd, 1 H, H-4), 7.7 (d, 1 H, H-3), 7.75 (dd, 1 H, H-6), 7.85 ("dd", 4 H, phenylene-H)

183.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-isopropylmercapto-phenyl)-benzamide

20



1.00 g (2.21 mmol) of 5-chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-mercapto-phenyl)-benzamide were dissolved in 25 ml of dimethylformamide and 0.25 g (2.21 mmol) of potassium tert-butyrate were added. The mixture was stirred at room temperature for 15 min, then 0.27 g (2.21 mmol) of isopropyl bromide were added dropwise, and the mixture was heated to 60 °C for 8 h. For working up it was poured onto water and extracted with ethyl acetate. The combined extracts were evaporated and the residue was purified by chromatography over silica with hexane/ethyl acetate (3:1). 420 mg (39 %) of the title compound having a melting point of 168-169 °C were obtained.

Analogously were obtained:

184.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-cyanomethylmercapto-phenyl)-benzamide; M.p.: 104 °C

185.) (4-((5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoyl)-amino)-phenylmercapto)-acetic acid ethyl ester; M.p.: 133 °C

186.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(2-(morpholin-4-yl)-ethylmercapto)-phenyl)-benzamide; M.p.: 95 °C

187.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(2-(2-methoxy-ethoxy)-ethylmercapto)-phenyl)-benzamide; Oil

188.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(prop-2-ynyl)-mercapto-phenyl)-benzamide; M.p.: 185 °C

189.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-isopropylmercapto-phenyl)-benzamide; M.p.: 169 °C

190.) 5-Chloro-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide sodium salt

A mixture of 0.48 g finely powdered sodium hydroxide and 7 g of 5-chloro-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide in 250 ml of ethanol was brought into solution by short heating. Then the mixture was evaporated in vacuo, 50 ml of water were added and it was again evaporated in vacuo to dryness. This procedure was repeated twice. The resulting product was dried in vacuo at 50 °C. M.p.: 343 °C (Dec.)

Analogously to the above compounds the following example compounds were obtained:

191.) 4,5-Dimethoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(N-methyl-N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide hydrochloride; M.p.: 214°C

192.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-3-methyl-phenyl)-benzamide; M.p.: 192 °C

193.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-3-methyl-phenyl)-benzamide; M.p.: 254 °C

194.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(3,5-dimethyl-piperidine-1-sulfonyl)-3-methyl-phenyl)-benzamide; M.p.: 242 °C

195.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(piperidine-1-sulfonyl)-3-methyl-phenyl)-benzamide; M.p.: 189 °C

196.) 4,5-Dimethoxy-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(N-methyl-N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; M.p.: 213 °C

197.) 4,5-Dimethoxy-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; M.p.: 216 °C

198.) 5-Chloro-2-(2,4-dimethyl-thiazole-5-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 190 °C

199.) 4,5-Dimethoxy-2-(4-chloro-phenylsulfonylamino)-N-(4-(3,5-dimethyl-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 249 °C (Dec.)

200.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(N-methyl-N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; Resin

201.) 3,4-Dimethoxy-2-(4-chloro-phenylsulfonylamino)-N-(4-(N-methyl-N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; M.p.: 241 °C

202.) 5-Bromo-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-3-methyl-phenyl)-benzamide; M.p.: 249 °C

203.) 5-Bromo-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 244 °C

204.) 5-Bromo-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 197 °C

205.) 4,5-Dimethoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(1,2,3,4-tetrahydro-isoquinoline-2-sulfonyl)-phenyl)-benzamide; M.p.: 213 °C

206.) 4,5-Dimethoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 232 °C

207.) 4,5-Dimethoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(cis-2,6-dimethyl-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 213 °C

208.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(1,2,3,4-tetrahydro-isoquinoline-2-sulfonyl)-phenyl)-benzamide; M.p.: 260 °C

209.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(N-methyl-N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; M.p.: 65 °C (sintering)

210.) 6-Methyl-2-(4-chloro-phenylsulfonylamino)-N-(4-(perhydroazepine-1-sulfonyl)-phenyl)-benzamide; M.p.: 151 °C

211.) 6-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(pyrrolidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 217 °C

212.) 6-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-hydroxybutylamino)-sulfonyl)-phenyl)-benzamide; Resin

213.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(N-ethyl-N-(pyridin-4-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; Resin

214.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 209 °C

215.) 3-Methyl-2-(4-chloro-phenylsulfonylamino)-N-(4-(N-methyl-N-(2-(pyridin-2-yl)-ethyl)-aminosulfonyl)-phenyl)-benzamide; M.p.: 193 °C

216.) 4,5-Difluoro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-aminocarbonyl-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 227 °C

217.) 4,5-Difluoro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-(2-hydroxyethyl)-piperazine-1-sulfonyl)-phenyl)-benzamide; Resin

218.) 5-Chloro-4-methoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; Oil

219.) 5-Chloro-4-methoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-1-sulfonyl)-phenyl)-benzamide; M.p.: 89 °C

220.) 5-Chloro-4-methoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(N-pyridin-3-yl)-N-methyl-aminosulfonyl)-phenyl)-benzamide; M.p.: 135 °C

221.) 4,5-Dimethoxy-2-(4-chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide sodium salt; M.p.: 330 °C (Dec)

222.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 230 °C

223.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(3,5-dimethylpiperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 61 °C

224.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 286 °C

225.) 5-Chloro-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-phenylsulfonyl)-phenyl-benzamide; M.p.: 227 °C

226.) 4-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 103 °C.

### Pharmacological investigations

#### (1) Activation of the soluble guanylate cyclase

The activation of the soluble guanylate cyclase (sGC) which catalyzes the conversion of guanosine triphosphate (GTP) into cyclic guanosine monophosphate (cGMP) and pyrophosphate, by the compounds according to the invention was quantified with the aid of an enzyme immunoassay (EIA) from Amersham. For this purpose the substances to be tested were initially incubated with sGC in microtiter plates, and the amount of the cGMP formed was then determined.

The sGC which was employed had been isolated from bovine lung (see Methods in Enzymology, Volume 195, p. 377). The test solutions (100 µl per well) contained 50 mM triethanolamine (TEA) buffer (pH 7.5), 3 mM MgCl<sub>2</sub>, 3 mM reduced glutathione (GSH), 0.1 mM GTP, 1 mM 3-isobutyl-1-methylxanthine (IBMX), suitably diluted enzyme solution and the substance to be tested or, in the control experiments, the solvent. The substances to be tested were dissolved in dimethyl sulfoxide (DMSO) and the solution was diluted with DMSO/water, so that the final concentration (c) of the substance to be tested in the test solution had the value indicated in the table. The DMSO concentration in the test solution was 5 % (v/v). The reaction was initiated by addition of the sGC. The

reaction mixture was incubated at 37 °C for 15 to 20 minutes and then stopped by ice-cooling and addition of the stop reagent (50 mM EDTA, pH 8.0). An aliquot of 50 µl was taken and used for determining the cGMP content using the acetylation protocol of the Amersham cGMP-EIA kit. The absorption of the samples was measured at 450 nm (reference wavelength 620 nm) in a microtiter plate reader. The cGMP concentration was determined using a standard curve which was obtained under the same test conditions. The activation of sGC by a test substance is given as the n-fold stimulation of the basal enzyme activity which was found in the control experiments (using solvent instead of test substance) (calculated using the formula

$$\text{n-fold stimulation} = [\text{cGMP}]_{\text{test substance}} / [\text{cGMP}]_{\text{control}}.$$

The following results were obtained:

Compound of	Concentration "c"	"n-fold" stimulation
Example No.	[µM]	
23	50	14.7
59	50	34.8
63	50	33.9
64	50	23.9
65	50	24.6
66	50	33
67	50	29.6
68	50	12.1
69	50	28.3
70	50	25.1
71	50	13.4
72	50	27
73	50	16.5
75	50	5.1
76	50	10.6
77	50	5.9
79	50	15.4
84	50	23.7



	Compound of	Concentration "c"	"n-fold" stimulation
	Example No.	[ $\mu$ M]	
	86	50	32.9
	87	50	12.5
	88	50	24.4
	89	50	11.6
5	124	50	31.2
	129	50	8.6
	130	50	35.3
	132	50	9.9
	134	50	7.2
10	136	50	24.2
	137	50	4.6
	139	50	21.9
	145	50	8
	146	50	10.2
15	148	50	15.5
	150	50	15.3
	151	50	19.9
	155	50	7.8
	156	50	7.8
20	157	50	4.6
	175	50	21.1
	176	50	13.9
	191	10	27.5
	192	10	26.7
25	193	10	31.1
	194	5	20.0
	195	10	16.2
	196	10	21.9
	206	50	19.3
30	208	10	23.1
	209	10	28.5
	222	10	29.5
	223	10	27.1
	224	50	27.8
35	225	10	13.4
	226	25	3.3

(2) Relaxation of rat aorta

For this test, normotensive male Wistar-Kyoto rats were sacrificed by a blow to the neck.

5 The abdominal cavity and the thorax were opened by a medium sternotomy. The descending aorta was subsequently removed, freed of connective tissue and divided into 8 rings of a length of approximately 4 mm. The tip of a pair of tweezers was introduced into the lumen of 4 of the 8 rings. The endothelium was removed by carefully rolling the rings over the tip of the pair of tweezers. All 8 aorta rings (4 with endothelium and 4 without endothelium) were subsequently suspended in an organ bath (Schuler-Organbad; Hugo Sachs Elektronik) at a constant temperature of 37 °C for the isometric determination of the contractile tone. For 30 minutes, the rings were calibrated at a resting tension of 1 g in carbonated (95 % O<sub>2</sub>; 5 % CO<sub>2</sub>) Krebs-Henseleit solution (composition: Na<sup>+</sup> 144.0 mM; K<sup>+</sup> 5.9 mM; Cl<sup>-</sup> 126.9 mM; Ca<sup>2+</sup> 1.6 mM; Mg<sup>2+</sup> 1.2 mM; H<sub>2</sub>PO<sub>4</sub><sup>-</sup> 1.2 mM; SO<sub>4</sub><sup>2-</sup> 1.2 mM; HCO<sub>3</sub><sup>-</sup> 25.0 mM; D-glucose 11.1 mM) of pH 7.4. Additionally, 1 µmol/l of indomethacin were added to the Krebs-Henseleit solution to inhibit prostaglandin biosynthesis. The rings were subsequently precontracted by addition of phenylephrine (concentration in the solution: 1 µM) and the endothelium-dependent relaxation or the functional loss of the endothelium was tested by addition of acetylcholine (concentration in the solution: 1 µM). After a 30-minute washing period, the rings were then again precontracted by addition of phenylephrine (1 µM), and the relaxing action of the test substances of the formula I was determined by administration of cumulative doses of the latter. The data were evaluated by standard methods. Given is the concentration IC<sub>50</sub> by which contraction is inhibited by 50 % (50 % relaxation).

25

The following results were obtained:

Compound of

IC<sub>50</sub>

Example No.

59	ring without endothelium	0.27 $\mu$ M
59	ring with endothelium	0.52 $\mu$ M
88	ring without endothelium	0.29 $\mu$ M
88	ring with endothelium	0.67 $\mu$ M
129	ring without endothelium	0.31 $\mu$ M
129	ring with endothelium	0.46 $\mu$ M

(3) Hemodynamic effect in the pig

Three pigs (German landrace) were anesthetized (Ketamine 20 mg/kg i.m., Methomidate 8 mg/kg i.p., Xylazine 2.5 mg/kg i.m. and Pentobarbital 25 mg/kg i.v. as a bolus plus 0.16 mg/kg per minute). The trachea was intubated and the animals were given artificial respiration with air. Oxygen was added to keep the blood gas parameters in the normal range. To record the blood pressure (BP; BP(s) = systolic blood pressure, BP(d) = diastolic blood pressure) by means of a Statham 23Db pressure transducer a catheter was inserted into the right A. Femoralis. The left ventricular pressure (LVP), the left ventricular end-diastolic pressure (LVEDP), the contractility (dP/dt) and the heart rate (HR) were determined with a Millar PC 350 catheter "tip manometer" which was inserted into the right ventricle. After a stabilization period of the hemodynamic parameters of 30 minutes the test substance was administered in the indicated dose into the exposed duodenum by means of a catheter. The determined data were evaluated according to standard methods. Given are the means and the standard deviations ( $M \pm SEM$ ) of the starting values and of the maximal changes of the individual parameters (= maximal effects).

The following results were obtained:

Compound of example 88 (dosage 10 mg/kg i.d.)

5	Parameter	starting value	maximal alteration	duration of action (min)
	BP(s) (mm Hg)	123 ± 26	-23 ± 6	> 180
	BP(d) (mm Hg)	83 ± 24	-20 ± 8	> 180
10	LVEDP (mm Hg)	4 ± 0.6	-1.3 ± 0.3	> 180
	dP/dt <sub>max</sub> (mm Hg/sec))	1800 ± 289	-633 ± 33	> 180
	HR (beats/min)	98 ± 2	-8 ± 2	> 180

The invention may be embodied in other specific forms and those skilled in the  
 art will recognize that various changes and modifications can be made without departing  
 from the spirit or essential characteristics thereof. The present embodiment is,  
 therefore, considered in all respects as illustrative and not restrictive, the scope of the  
 appended claims rather than by the foregoing description, and all changes which come  
 within the meaning and range of equivalency of the claims are therefore intended to be  
 embraced therein.